

L^p -cohomology of symmetric spaces

P. Pansu

September 29, 2006

What is it ?

topological space \rightarrow *cohomology*
manifold \rightarrow *de Rham cohomology*

What is it ?

| | | |
|----------------------------|---|--|
| <i>topological space</i> | → | <i>cohomology</i> |
| <i>manifold</i> | → | <i>de Rham cohomology</i> |
| <i>metric space</i> | → | <i>cohomology with decay condition</i> |
| <i>Riemannian manifold</i> | → | <i>de Rham cohomology with decay condition</i> |

What is it ?

| | | |
|----------------------------|---|--|
| <i>topological space</i> | → | <i>cohomology</i> |
| <i>manifold</i> | → | <i>de Rham cohomology</i> |
| <i>metric space</i> | → | <i>cohomology with decay condition</i> |
| <i>Riemannian manifold</i> | → | <i>de Rham cohomology with decay condition</i> |

Definition

Let M be a Riemannian manifold. Let $p > 1$. L^p -cohomology of M is the cohomology of the complex of L^p -differential forms on M whose exterior differentials are L^p as well,

$$H^{k,p} = \text{closed } k\text{-forms in } L^p / d((k-1)\text{-forms in } L^p),$$

What is it ?

| | | |
|----------------------------|---|--|
| <i>topological space</i> | → | <i>cohomology</i> |
| <i>manifold</i> | → | <i>de Rham cohomology</i> |
| <i>metric space</i> | → | <i>cohomology with decay condition</i> |
| <i>Riemannian manifold</i> | → | <i>de Rham cohomology with decay condition</i> |

Definition

Let M be a Riemannian manifold. Let $p > 1$. L^p -cohomology of M is the cohomology of the complex of L^p -differential forms on M whose exterior differentials are L^p as well,

$$H^{k,p} = \text{closed } k\text{-forms in } L^p / d((k-1)\text{-forms in } L^p),$$

$$R^{k,p} = \text{closed } k\text{-forms in } L^p / \text{closure of } d((k-1)\text{-forms in } L^p),$$

$$T^{k,p} = \text{closure of } d((k-1)\text{-forms in } L^p) / d((k-1)\text{-forms in } L^p).$$

$R^{k,p}$ is called the reduced cohomology. $T^{k,p}$ is called the torsion.

Example : the real hyperbolic plane $H_{\mathbb{R}}^2$

Here $H^{0,p} = 0 = H^{2,p}$ for all p .

Example : the real hyperbolic plane $H_{\mathbb{R}}^2$

Here $H^{0,p} = 0 = H^{2,p}$ for all p .

If $p = 2$, since the Laplacian on L^2 functions is bounded below, $T^{1,2} = 0$. Therefore

$$H^{1,2} = R^{1,2}$$

Example : the real hyperbolic plane $H_{\mathbb{R}}^2$

Here $H^{0,p} = 0 = H^{2,p}$ for all p .

If $p = 2$, since the Laplacian on L^2 functions is bounded below, $T^{1,2} = 0$. Therefore

$$\begin{aligned} H^{1,2} &= R^{1,2} \\ &= \{L^2 \text{ harmonic 1-forms}\} \end{aligned}$$

Example : the real hyperbolic plane $H_{\mathbb{R}}^2$

Here $H^{0,p} = 0 = H^{2,p}$ for all p .

If $p = 2$, since the Laplacian on L^2 functions is bounded below, $T^{1,2} = 0$. Therefore

$$\begin{aligned} H^{1,2} &= R^{1,2} \\ &= \{L^2 \text{ harmonic 1-forms}\} \\ &= \{\text{harmonic functions } h \text{ on } H_{\mathbb{R}}^2 \text{ with } \nabla h \in L^2\} / \mathbb{R}. \end{aligned}$$

Example : the real hyperbolic plane $H_{\mathbb{R}}^2$

Here $H^{0,p} = 0 = H^{2,p}$ for all p .

If $p = 2$, since the Laplacian on L^2 functions is bounded below, $T^{1,2} = 0$. Therefore

$$\begin{aligned} H^{1,2} &= R^{1,2} \\ &= \{L^2 \text{ harmonic 1-forms}\} \\ &= \{\text{harmonic functions } h \text{ on } H_{\mathbb{R}}^2 \text{ with } \nabla h \in L^2\} / \mathbb{R}. \end{aligned}$$

Using conformal invariance, switch from hyperbolic metric to euclidean metric on the disk D .

$$H^{1,2} = \{\text{harmonic functions } h \text{ on } D \text{ with } \nabla h \in L^2\} / \mathbb{R}$$

Example : the real hyperbolic plane $H_{\mathbb{R}}^2$

Here $H^{0,p} = 0 = H^{2,p}$ for all p .

If $p = 2$, since the Laplacian on L^2 functions is bounded below, $T^{1,2} = 0$. Therefore

$$\begin{aligned} H^{1,2} &= R^{1,2} \\ &= \{L^2 \text{ harmonic 1-forms}\} \\ &= \{\text{harmonic functions } h \text{ on } H_{\mathbb{R}}^2 \text{ with } \nabla h \in L^2\} / \mathbb{R}. \end{aligned}$$

Using conformal invariance, switch from hyperbolic metric to euclidean metric on the disk D .

$$\begin{aligned} H^{1,2} &= \{\text{harmonic functions } h \text{ on } D \text{ with } \nabla h \in L^2\} / \mathbb{R} \\ &= \{\text{Fourier series } \sum a_n e^{in\theta} \text{ with } a_0 = 0, \sum |n| |a_n|^2 < +\infty\}, \end{aligned}$$

Example : the real hyperbolic plane $H_{\mathbb{R}}^2$

Here $H^{0,p} = 0 = H^{2,p}$ for all p .

If $p = 2$, since the Laplacian on L^2 functions is bounded below, $T^{1,2} = 0$. Therefore

$$\begin{aligned} H^{1,2} &= R^{1,2} \\ &= \{L^2 \text{ harmonic 1-forms}\} \\ &= \{\text{harmonic functions } h \text{ on } H_{\mathbb{R}}^2 \text{ with } \nabla h \in L^2\} / \mathbb{R}. \end{aligned}$$

Using conformal invariance, switch from hyperbolic metric to euclidean metric on the disk D .

$$\begin{aligned} H^{1,2} &= \{\text{harmonic functions } h \text{ on } D \text{ with } \nabla h \in L^2\} / \mathbb{R} \\ &= \{\text{Fourier series } \sum a_n e^{in\theta} \text{ with } a_0 = 0, \sum |n| |a_n|^2 < +\infty\}, \end{aligned}$$

which is Sobolev space $H^{1/2}(\mathbb{R}/2\pi\mathbb{Z})$ mod constants.

Example : the real hyperbolic plane $H_{\mathbb{R}}^2$

Here $H^{0,p} = 0 = H^{2,p}$ for all p .

If $p = 2$, since the Laplacian on L^2 functions is bounded below, $T^{1,2} = 0$. Therefore

$$\begin{aligned} H^{1,2} &= R^{1,2} \\ &= \{L^2 \text{ harmonic 1-forms}\} \\ &= \{\text{harmonic functions } h \text{ on } H_{\mathbb{R}}^2 \text{ with } \nabla h \in L^2\} / \mathbb{R}. \end{aligned}$$

Using conformal invariance, switch from hyperbolic metric to euclidean metric on the disk D .

$$\begin{aligned} H^{1,2} &= \{\text{harmonic functions } h \text{ on } D \text{ with } \nabla h \in L^2\} / \mathbb{R} \\ &= \{\text{Fourier series } \sum a_n e^{in\theta} \text{ with } a_0 = 0, \sum |n| |a_n|^2 < +\infty\}, \end{aligned}$$

which is Sobolev space $H^{1/2}(\mathbb{R}/2\pi\mathbb{Z})$ mod constants.

More generally, for $p > 1$, $T^{1,p} = 0$ and $H^{1,p}$ is equal to the Besov space $B_{p,p}^{1/p}(\mathbb{R}/2\pi\mathbb{Z})$ mod constants.

$$H^{0,p} = 0.$$

$$H^{0,p} = 0.$$

$$R^{1,p} = 0,$$

$$H^{0,p} = 0.$$

$$R^{1,p} = 0,$$

since every function in $L^p(\mathbb{R})$ can be approximated in L^p with derivatives of compactly supported functions. Therefore $H^{1,p}$ is only torsion.

$$H^{0,p} = 0.$$

$$R^{1,p} = 0,$$

since every function in $L^p(\mathbb{R})$ can be approximated in L^p with derivatives of compactly supported functions. Therefore $H^{1,p}$ is only torsion.

$T^{1,p}$ is non zero and thus infinite dimensional.

$$H^{0,p} = 0.$$

$$R^{1,p} = 0,$$

since every function in $L^p(\mathbb{R})$ can be approximated in L^p with derivatives of compactly supported functions. Therefore $H^{1,p}$ is only torsion.

$T^{1,p}$ is non zero and thus infinite dimensional.

Indeed, the 1-form $\frac{dt}{t}$ (cut off near the origin) is in L^p for all $p > 1$ but it is not the differential of a function in L^p .

What are our favourite spaces ?

- ▶ L^p -cohomology has been used (L. Saper, S. Zucker) to study manifolds with **thin ends**. The answer is related to the topology of a compactification.
- ▶ In this talk : manifolds with **large ends**, e.g. symmetric spaces themselves. L^p -cohomology is related to analytic features of a compactification.

cohomology \rightarrow continuous maps

cohomology \rightarrow continuous maps
 L^p -cohomology \rightarrow uniform maps.

cohomology \rightarrow continuous maps
 L^p -cohomology \rightarrow uniform maps.

Definition

A map $f : X \rightarrow Y$ between metric spaces is uniform if $d(f(x), f(x'))$ is bounded from above in terms of $d(x, x')$ only.

cohomology \rightarrow continuous maps
 L^p -cohomology \rightarrow uniform maps.

Definition

A map $f : X \rightarrow Y$ between metric spaces is uniform if $d(f(x), f(x'))$ is bounded from above in terms of $d(x, x')$ only.

Examples

The obvious map $\mathbb{Z} \rightarrow \mathbb{R}$ is uniform. Any homomorphism between groups (with left invariant metrics) is uniform. The parametrization of a cusp by a punctured disk is not uniform.


cohomology \rightarrow continuous maps
 L^p -cohomology \rightarrow uniform maps.

Definition

A map $f : X \rightarrow Y$ between metric spaces is uniform if $d(f(x), f(x'))$ is bounded from above in terms of $d(x, x')$ only.

Examples

The obvious map $\mathbb{Z} \rightarrow \mathbb{R}$ is uniform. Any homomorphism between groups (with left invariant metrics) is uniform. The parametrization of a cusp by a punctured disk is not uniform.



Proposition

Among contractible Riemannian manifolds admitting a cocompact isometric group action, L^p -cohomology is natural under uniform maps.

L^p -cohomology can be discretized.

L^p -cohomology can be discretized. It makes sense for discrete groups, and cannot see any difference between a cocompact lattice in a semi-simple Lie group G , the Lie group G itself or the Riemannian symmetric space G/K .

L^p -cohomology can be discretized. It makes sense for discrete groups, and cannot see any difference between a cocompact lattice in a semi-simple Lie group G , the Lie group G itself or the Riemannian symmetric space G/K .

In conclusion,

- ▶ L^p -cohomology is a tool to investigate discrete groups.
- ▶ It shares nearly all properties of usual cohomology.
- ▶ Nevertheless, it is not easy to calculate it.
- ▶ In the case of cocompact lattices in Lie groups, it can probably be computed by analytic means.

L^p -cohomology can be discretized. It makes sense for discrete groups, and cannot see any difference between a cocompact lattice in a semi-simple Lie group G , the Lie group G itself or the Riemannian symmetric space G/K .

In conclusion,

- ▶ L^p -cohomology is a tool to investigate discrete groups.
- ▶ It shares nearly all properties of usual cohomology.
- ▶ Nevertheless, it is not easy to calculate it.
- ▶ In the case of cocompact lattices in Lie groups, it can probably be computed by analytic means.

In this talk, we explain 3 applications of L^p -cohomology to negatively curved Riemannian manifolds and groups.

1. Hopf's conjecture about Euler characteristic
2. Cannon conjecture on groups with boundary a 2-sphere
3. Curvature pinching

1. Hopf's conjecture about Euler characteristic
2. Cannon conjecture on groups with boundary a 2-sphere
3. Curvature pinching

Remark

- ▶ *Compact 2-dimensional negatively curved manifolds have negative Euler characteristic.*
- ▶ *2m-dimensional compact hyperbolic manifolds have Euler characteristic proportional to $(-1)^m$.*
- ▶ *This generalizes to all compact negatively curved locally symmetric spaces (Gauss-Bonnet).*

Remark

- ▶ *Compact 2-dimensional negatively curved manifolds have negative Euler characteristic.*
- ▶ *2m-dimensional compact hyperbolic manifolds have Euler characteristic proportional to $(-1)^m$.*
- ▶ *This generalizes to all compact negatively curved locally symmetric spaces (Gauss-Bonnet).*

Conjecture

(H. Hopf). If M is 2m-dimensional compact negatively curved, then $(-1)^m \chi(M) > 0$.

Remark

- ▶ *Compact 2-dimensional negatively curved manifolds have negative Euler characteristic.*
- ▶ *2m-dimensional compact hyperbolic manifolds have Euler characteristic proportional to $(-1)^m$.*
- ▶ *This generalizes to all compact negatively curved locally symmetric spaces (Gauss-Bonnet).*

Conjecture

(H. Hopf). If M is 2m-dimensional compact negatively curved, then $(-1)^m \chi(M) > 0$.

Theorem

(M. Gromov, 1991). This is true provided M also admits a Kähler metric.

Definition

A Riemannian manifold M is Kähler if it admits a parallel complex structure.

Definition

A Riemannian manifold M is Kähler if it admits a parallel complex structure.

Then M is a complex manifold. Every complex submanifold in complex projective space admits a Kähler metric.

Definition

A Riemannian manifold M is Kähler if it admits a parallel complex structure.

Then M is a complex manifold. Every complex submanifold in complex projective space admits a Kähler metric.

Proposition

(Part of hard Lefschetz theorem). Let M^{2m} be a compact Kähler manifold with Kähler form ω . Then wedging with ω maps harmonic forms to harmonic forms,

Definition

A Riemannian manifold M is Kähler if it admits a parallel complex structure.

Then M is a complex manifold. Every complex submanifold in complex projective space admits a Kähler metric.

Proposition

(Part of hard Lefschetz theorem). Let M^{2m} be a compact Kähler manifold with Kähler form ω . Then wedging with ω maps harmonic forms to harmonic forms, and this induces an injection in cohomology $H^k(M, \mathbb{R}) \rightarrow H^{k+2}(M, \mathbb{R})$ for all $k < m$.

Definition

A Riemannian manifold M is Kähler if it admits a parallel complex structure.

Then M is a complex manifold. Every complex submanifold in complex projective space admits a Kähler metric.

Proposition

(Part of hard Lefschetz theorem). Let M^{2m} be a compact Kähler manifold with Kähler form ω . Then wedging with ω maps harmonic forms to harmonic forms, and this induces an injection in cohomology $H^k(M, \mathbb{R}) \rightarrow H^{k+2}(M, \mathbb{R})$ for all $k < m$.

Corollary

Let M^{2m} be a complete Kähler manifold with Kähler form ω . Then wedging with ω maps L^2 -harmonic forms to L^2 -harmonic forms, and this induces an injection in reduced L^2 -cohomology $R^{k,2}(M) \rightarrow R^{k+2}(M)$ for all $k < m$.

Proposition

(M. Gromov). Let \tilde{M} be a complete simply connected negatively curved Riemannian manifold. Let $k \geq 2$.

- ▶ *(Coning of cycles). Every $k - 1$ -cycle z spans a k -chain c with $\text{vol}(c) \leq \text{const. vol}(z)$.*

Proposition

(M. Gromov). Let \tilde{M} be a complete simply connected negatively curved Riemannian manifold. Let $k \geq 2$.

- ▶ *(Coning of cycles). Every $k - 1$ -cycle z spans a k -chain c with $\text{vol}(c) \leq \text{const. vol}(z)$.*
- ▶ *(Coning of forms). Every closed bounded differential k -form α on \tilde{M} is the differential of a bounded $(k - 1)$ -form β with $\|\beta\|_{L^\infty} \leq \text{const.} \|\alpha\|_{L^\infty}$.*

Proposition

(M. Gromov). Let \tilde{M} be a complete simply connected negatively curved Riemannian manifold. Let $k \geq 2$.

- ▶ *(Coning of cycles). Every $k - 1$ -cycle z spans a k -chain c with $\text{vol}(c) \leq \text{const. vol}(z)$.*
- ▶ *(Coning of forms). Every closed bounded differential k -form α on \tilde{M} is the differential of a bounded $(k - 1)$ -form β with $\|\beta\|_{L^\infty} \leq \text{const.} \|\alpha\|_{L^\infty}$.*

Corollary

Assume \tilde{M}^{2m} covers both a compact Kähler manifold and a compact negatively curved Riemannian manifold. Then $R^{k,2}(\tilde{M}) = 0$ for all $k \neq m$.

Proposition

(M. Gromov). Let \tilde{M} be a complete simply connected negatively curved Riemannian manifold. Let $k \geq 2$.

- ▶ *(Coning of cycles). Every $k - 1$ -cycle z spans a k -chain c with $\text{vol}(c) \leq \text{const. vol}(z)$.*
- ▶ *(Coning of forms). Every closed bounded differential k -form α on \tilde{M} is the differential of a bounded $(k - 1)$ -form β with $\|\beta\|_{L^\infty} \leq \text{const.} \|\alpha\|_{L^\infty}$.*

Corollary

Assume \tilde{M}^{2m} covers both a compact Kähler manifold and a compact negatively curved Riemannian manifold. Then $R^{k,2}(\tilde{M}) = 0$ for all $k \neq m$. Furthermore, $T^{,2}(\tilde{M}) = 0$.*

Proposition

(M. Gromov). Let \tilde{M} be a complete simply connected negatively curved Riemannian manifold. Let $k \geq 2$.

- ▶ *(Coning of cycles). Every $k - 1$ -cycle z spans a k -chain c with $\text{vol}(c) \leq \text{const. vol}(z)$.*
- ▶ *(Coning of forms). Every closed bounded differential k -form α on \tilde{M} is the differential of a bounded $(k - 1)$ -form β with $\|\beta\|_{L^\infty} \leq \text{const.} \|\alpha\|_{L^\infty}$.*

Corollary

Assume \tilde{M}^{2m} covers both a compact Kähler manifold and a compact negatively curved Riemannian manifold. Then $R^{k,2}(\tilde{M}) = 0$ for all $k \neq m$. Furthermore, $T^{,2}(\tilde{M}) = 0$.*

Proof. Lift Kähler form to universal cover \tilde{M} . Write $\omega = db$ with b bounded.

Proposition

(M. Gromov). Let \tilde{M} be a complete simply connected negatively curved Riemannian manifold. Let $k \geq 2$.

- ▶ *(Coning of cycles). Every $k - 1$ -cycle z spans a k -chain c with $\text{vol}(c) \leq \text{const. vol}(z)$.*
- ▶ *(Coning of forms). Every closed bounded differential k -form α on \tilde{M} is the differential of a bounded $(k - 1)$ -form β with $\|\beta\|_{L^\infty} \leq \text{const.} \|\alpha\|_{L^\infty}$.*

Corollary

Assume \tilde{M}^{2m} covers both a compact Kähler manifold and a compact negatively curved Riemannian manifold. Then $R^{k,2}(\tilde{M}) = 0$ for all $k \neq m$. Furthermore, $T^{,2}(\tilde{M}) = 0$.*

Proof. Lift Kähler form to universal cover \tilde{M} . Write $\omega = db$ with b bounded. Let $k < m$. For α a closed k -form in L^2 ,

$$\omega \wedge \alpha = d(b \wedge \alpha) \quad \text{and} \quad b \wedge \alpha \in L^2,$$

Proposition

(M. Gromov). Let \tilde{M} be a complete simply connected negatively curved Riemannian manifold. Let $k \geq 2$.

- ▶ *(Coning of cycles). Every $k - 1$ -cycle z spans a k -chain c with $\text{vol}(c) \leq \text{const. vol}(z)$.*
- ▶ *(Coning of forms). Every closed bounded differential k -form α on \tilde{M} is the differential of a bounded $(k - 1)$ -form β with $\|\beta\|_{L^\infty} \leq \text{const.} \|\alpha\|_{L^\infty}$.*

Corollary

Assume \tilde{M}^{2m} covers both a compact Kähler manifold and a compact negatively curved Riemannian manifold. Then $R^{k,2}(\tilde{M}) = 0$ for all $k \neq m$. Furthermore, $T^{,2}(\tilde{M}) = 0$.*

Proof. Lift Kähler form to universal cover \tilde{M} . Write $\omega = db$ with b bounded. Let $k < m$. For α a closed k -form in L^2 ,

$$\omega \wedge \alpha = d(b \wedge \alpha) \quad \text{and} \quad b \wedge \alpha \in L^2,$$

thus $\omega \wedge \alpha = 0$ in $R^{k+2,2}(\tilde{M})$. If α is harmonic, conclude that $\alpha = 0$ in $R^{k,2}(\tilde{M})$.

Let \tilde{M} cover a compact manifold M . If nonzero, $R^{k,2}(\tilde{M})$ is infinite dimensional.

Let \tilde{M} cover a compact manifold M . If nonzero, $R^{k,2}(\tilde{M})$ is infinite dimensional. Nevertheless, **M. Atiyah** defined a *von Neumann dimension*

$$b^{k,2}(M) = \dim_{vN} R^{k,2}(\tilde{M}),$$

called the k -th L^2 -Betti number of M .

Let \tilde{M} cover a compact manifold M . If nonzero, $R^{k,2}(\tilde{M})$ is infinite dimensional. Nevertheless, **M. Atiyah** defined a *von Neumann dimension*

$$b^{k,2}(M) = \dim_{vN} R^{k,2}(\tilde{M}),$$

called the k -th L^2 -Betti number of M .

Examples

(W. Lück). If M admits a tower of finite degree d_j normal coverings M_j such that $\bigcap_j \pi_1(M_j) = \{1\}$, then

$$b^{k,2}(M) = \lim_{j \rightarrow \infty} \frac{b^k(M_j, \mathbb{R})}{d_j}.$$

Let \tilde{M} cover a compact manifold M . If nonzero, $R^{k,2}(\tilde{M})$ is infinite dimensional. Nevertheless, **M. Atiyah** defined a *von Neumann dimension*

$$b^{k,2}(M) = \dim_{vN} R^{k,2}(\tilde{M}),$$

called the k -th L^2 -Betti number of M .

Examples

(W. Lück). If M admits a tower of finite degree d_j normal coverings M_j such that $\bigcap_j \pi_1(M_j) = \{1\}$, then

$$b^{k,2}(M) = \lim_{j \rightarrow \infty} \frac{b^k(M_j, \mathbb{R})}{d_j}.$$

Proposition

Let \tilde{M} cover a compact manifold M . Then

$$\chi(M) = \sum_k (-1)^k b^{k,2}(M).$$

Proposition

(Relative index theorem, M. Gromov-B. Lawson). Let \tilde{M} be a simply connected nonpositively curved Riemannian manifold. Then there exists k such that $H^{k,2}(\tilde{M}) \neq 0$.

Proposition

(Relative index theorem, M. Gromov-B. Lawson). Let \tilde{M} be a simply connected nonpositively curved Riemannian manifold. Then there exists k such that $H^{k,2}(\tilde{M}) \neq 0$.

Proof of Gromov's theorem. Assume M is compact and admits both a negatively curved metric and a Kähler metric. Then all $b^{k,2}(M)$ vanish except $b^{m,2}(M)$, which is nonzero, thus $(-1)^m \chi(M) = b^{m,2}(M) > 0$.

Proposition

(Relative index theorem, M. Gromov-B. Lawson). Let \tilde{M} be a simply connected nonpositively curved Riemannian manifold. Then there exists k such that $H^{k,2}(\tilde{M}) \neq 0$.

Proof of Gromov's theorem. Assume M is compact and admits both a negatively curved metric and a Kähler metric. Then all $b^{k,2}(M)$ vanish except $b^{m,2}(M)$, which is nonzero, thus $(-1)^m \chi(M) = b^{m,2}(M) > 0$.

In conclusion, we have used

- ▶ Lefschetz mechanism, L^2 -Betti numbers.

Proposition

(Relative index theorem, M. Gromov-B. Lawson). Let \tilde{M} be a simply connected nonpositively curved Riemannian manifold. Then there exists k such that $H^{k,2}(\tilde{M}) \neq 0$.

Proof of Gromov's theorem. Assume M is compact and admits both a negatively curved metric and a Kähler metric. Then all $b^{k,2}(M)$ vanish except $b^{m,2}(M)$, which is nonzero, thus $(-1)^m \chi(M) = b^{m,2}(M) > 0$.

In conclusion, we have used

- ▶ Lefschetz mechanism, L^2 -Betti numbers.
- ▶ Vanishing of L^∞ -cohomology.

Proposition

(Relative index theorem, M. Gromov-B. Lawson). Let \tilde{M} be a simply connected nonpositively curved Riemannian manifold. Then there exists k such that $H^{k,2}(\tilde{M}) \neq 0$.

Proof of Gromov's theorem. Assume M is compact and admits both a negatively curved metric and a Kähler metric. Then all $b^{k,2}(M)$ vanish except $b^{m,2}(M)$, which is nonzero, thus $(-1)^m \chi(M) = b^{m,2}(M) > 0$.

In conclusion, we have used

- ▶ Lefschetz mechanism, L^2 -Betti numbers.
- ▶ Vanishing of L^∞ -cohomology.
- ▶ Cup-product $H^{k,2} \otimes H^{2,\infty} \rightarrow H^{k+2,\infty}$.

1. Hopf's conjecture about Euler characteristic
2. Cannon conjecture on groups with boundary a 2-sphere
3. Curvature pinching

Remark

In negatively curved manifolds, triangles are thin.

Remark

In negatively curved manifolds, triangles are thin.

Definition

(M. Gromov 1986). Say a finitely generated group is hyperbolic if its triangles are thin.

Remark

In negatively curved manifolds, triangles are thin.

Definition

(M. Gromov 1986). Say a finitely generated group is hyperbolic if its triangles are thin. A hyperbolic group has a functorial ideal boundary.

Remark

In negatively curved manifolds, triangles are thin.

Definition

(M. Gromov 1986). Say a finitely generated group is hyperbolic if its triangles are thin. A hyperbolic group has a functorial ideal boundary.

Examples

- ▶ *A free group is hyperbolic, its ideal boundary is totally disconnected.*

Remark

In negatively curved manifolds, triangles are thin.

Definition

(M. Gromov 1986). Say a finitely generated group is hyperbolic if its triangles are thin. A hyperbolic group has a functorial ideal boundary.

Examples

- ▶ *A free group is hyperbolic, its ideal boundary is totally disconnected.*
- ▶ *The fundamental group of a compact negatively curved n -manifold is hyperbolic, its ideal boundary is an $n - 1$ -sphere.*

Remark

In negatively curved manifolds, triangles are thin.

Definition

(M. Gromov 1986). Say a finitely generated group is hyperbolic if its triangles are thin. A hyperbolic group has a functorial ideal boundary.

Examples

- ▶ *A free group is hyperbolic, its ideal boundary is totally disconnected.*
- ▶ *The fundamental group of a compact negatively curved n -manifold is hyperbolic, its ideal boundary is an $n - 1$ -sphere.*

Conjecture

(J. Cannon). Let Γ be a hyperbolic group whose ideal boundary is a 2-sphere. Then Γ is virtually a cocompact lattice in $PSL(2, \mathbb{C})$.

Remark

The ideal boundary of a hyperbolic group carries a natural conformal structure, but no canonical metric.

Remark

The ideal boundary of a hyperbolic group carries a natural conformal structure, but no canonical metric.

Definition

Define the conformal dimension of a hyperbolic group as the least Hausdorff dimension of a metric in the natural conformal structure.

Remark

The ideal boundary of a hyperbolic group carries a natural conformal structure, but no canonical metric.

Definition

Define the conformal dimension of a hyperbolic group as the least Hausdorff dimension of a metric in the natural conformal structure.

Strategy (B. Kleiner). Prove that this infimum is achieved. Then prove that dimension minimizing metrics are Riemannian if boundary is 2-dimensional. Then apply a result of **D. Sullivan (1978)**: every uniformly quasiconformal group of the standard 2-sphere is conjugate to a subgroup of $PSL(2, \mathbb{C})$.

Remark

The ideal boundary of a hyperbolic group carries a natural conformal structure, but no canonical metric.

Definition

Define the conformal dimension of a hyperbolic group as the least Hausdorff dimension of a metric in the natural conformal structure.

Strategy (B. Kleiner). Prove that this infimum is achieved. Then prove that dimension minimizing metrics are Riemannian if boundary is 2-dimensional. Then apply a result of **D. Sullivan (1978)**: every uniformly quasiconformal group of the standard 2-sphere is conjugate to a subgroup of $PSL(2, \mathbb{C})$.

Theorem

(S. Keith-T. Laakso, M. Bonk-B. Kleiner 2005). Let Γ be a hyperbolic group whose ideal boundary is a 2-sphere. If conformal dimension is achieved, then Γ is virtually a cocompact lattice in $PSL(2, \mathbb{C})$.

Remark

For hyperbolic groups, $H^{1,p}$ is nonzero for p large, but zero for p small.

Remark

For hyperbolic groups, $H^{1,p}$ is nonzero for p large, but zero for p small. For instance, $H^{1,p}(SO(n, 1)) = 0 \Leftrightarrow p \leq n - 1$.

Definition

Define the L^p -dimension of a group as the least $p > 1$ such that its $H^{1,p}$ is nonzero.

Remark

For hyperbolic groups, $H^{1,p}$ is nonzero for p large, but zero for p small. For instance, $H^{1,p}(SO(n, 1)) = 0 \Leftrightarrow p \leq n - 1$.

Definition

Define the L^p -dimension of a group as the least $p > 1$ such that its $H^{1,p}$ is nonzero.

Theorem

(Same people + M. Bourdon-H. Pajot 2003). Let Γ be a hyperbolic group. Then L^p -dimension is less than or equal to conformal dimension.

Remark

For hyperbolic groups, $H^{1,p}$ is nonzero for p large, but zero for p small. For instance, $H^{1,p}(SO(n, 1)) = 0 \Leftrightarrow p \leq n - 1$.

Definition

Define the L^p -dimension of a group as the least $p > 1$ such that its $H^{1,p}$ is nonzero.

Theorem

(Same people + M. Bourdon-H. Pajot 2003). Let Γ be a hyperbolic group. Then L^p -dimension is less than or equal to conformal dimension. If conformal dimension is achieved, then L^p -dimension and conformal dimension coincide.

Examples

(M. Bourdon-H. Pajot). There exist hyperbolic groups for which conformal dimension $> 2 \geq L^p$ -dimension. For such groups, conformal dimension cannot be achieved.

In conclusion, we have used

- ▶ Mayer-Vietoris and L^2 -Betti numbers.
- ▶ Expression of $H^{1,p}$ as a function space on the ideal boundary.

1. Hopf's conjecture about Euler characteristic
2. Cannon conjecture on groups with boundary a 2-sphere
3. **Curvature pinching**

Remark

Rank one symmetric spaces are hyperbolic spaces over the reals $H_{\mathbb{R}}^n$, the complex numbers $H_{\mathbb{C}}^m$, the quaternions $H_{\mathbb{H}}^m$, and the octonions $H_{\mathbb{O}}^2$.

Real hyperbolic space has sectional curvature -1 . Other rank one symmetric spaces are $-\frac{1}{4}$ -pinched, i.e. their sectional curvature ranges between -1 and $-\frac{1}{4}$.

Remark

Rank one symmetric spaces are hyperbolic spaces over the reals $H_{\mathbb{R}}^n$, the complex numbers $H_{\mathbb{C}}^m$, the quaternions $H_{\mathbb{H}}^m$, and the octonions $H_{\mathbb{O}}^2$.

Real hyperbolic space has sectional curvature -1 . Other rank one symmetric spaces are $-\frac{1}{4}$ -pinched, i.e. their sectional curvature ranges between -1 and $-\frac{1}{4}$.

Definition

Define the optimal pinching $\delta(G)$ of a discrete (or Lie) group G as the least $\delta > -1$ such that G is bi-uniformly equivalent to a δ -pinched Riemannian manifold.

Remark

Rank one symmetric spaces are hyperbolic spaces over the reals $H_{\mathbb{R}}^n$, the complex numbers $H_{\mathbb{C}}^m$, the quaternions $H_{\mathbb{H}}^m$, and the octonions $H_{\mathbb{O}}^2$.

Real hyperbolic space has sectional curvature -1 . Other rank one symmetric spaces are $-\frac{1}{4}$ -pinched, i.e. their sectional curvature ranges between -1 and $-\frac{1}{4}$.


Definition

Define the optimal pinching $\delta(G)$ of a discrete (or Lie) group G as the least $\delta > -1$ such that G is bi-uniformly equivalent to a δ -pinched Riemannian manifold.

Conjecture

The optimal pinching of $SU(m, 1)$, $Sp(m, 1)$ ($m \geq 2$) and F_4^{-20} is $-\frac{1}{4}$.

L^p -cohomology of $H_{\mathbb{R}}^n$



 non vanishing reduced cohomology

 non vanishing torsion

Theorem

If M^n is simply connected and δ -pinched for some $\delta \in [-1, 0)$, then

$$p < 1 + \frac{n-k}{k-1} \sqrt{-\delta} \quad \Rightarrow \quad T^{k,p}(M) = 0.$$

Theorem

If M^n is simply connected and δ -pinched for some $\delta \in [-1, 0)$, then

$$p < 1 + \frac{n-k}{k-1} \sqrt{-\delta} \quad \Rightarrow \quad T^{k,p}(M) = 0.$$

This is sharp. For instance, consider the semidirect product $G = \mathbb{R}^3 \rtimes_{\alpha} \mathbb{R}$ where $\alpha = \text{diag}(1, 1, 2)$.

- ▶ It admits a $-\frac{1}{4}$ -pinched left-invariant Riemannian metric, therefore $\delta(G) \leq -\frac{1}{4}$.
- ▶ It has $T^{2,p}(G) \neq 0$ for $2 < p \leq 4$. This implies that $\delta(G) = -\frac{1}{4}$.

Theorem

If M^n is simply connected and δ -pinched for some $\delta \in [-1, 0)$, then

$$p < 1 + \frac{n-k}{k-1} \sqrt{-\delta} \quad \Rightarrow \quad T^{k,p}(M) = 0.$$

This is sharp. For instance, consider the semidirect product $G = \mathbb{R}^3 \rtimes_{\alpha} \mathbb{R}$ where $\alpha = \text{diag}(1, 1, 2)$.

- ▶ It admits a $-\frac{1}{4}$ -pinched left-invariant Riemannian metric, therefore $\delta(G) \leq -\frac{1}{4}$.
- ▶ It has $T^{2,p}(G) \neq 0$ for $2 < p \leq 4$. This implies that $\delta(G) = -\frac{1}{4}$.

Remark

Complex hyperbolic plane $H_{\mathbb{C}}^2$ is isometric to $G' = \text{Heis}^3 \rtimes_{\alpha} \mathbb{R}$ where $\alpha = \text{diag}(1, 1, 2)$ and Heis denotes the Heisenberg group. Therefore it is very close to G .

Theorem

If M^n is simply connected and δ -pinched for some $\delta \in [-1, 0)$, then

$$p < 1 + \frac{n-k}{k-1} \sqrt{-\delta} \quad \Rightarrow \quad T^{k,p}(M) = 0.$$

This is sharp. For instance, consider the semidirect product $G = \mathbb{R}^3 \rtimes_{\alpha} \mathbb{R}$ where $\alpha = \text{diag}(1, 1, 2)$.

- ▶ It admits a $-\frac{1}{4}$ -pinched left-invariant Riemannian metric, therefore $\delta(G) \leq -\frac{1}{4}$.
- ▶ It has $T^{2,p}(G) \neq 0$ for $2 < p \leq 4$. This implies that $\delta(G) = -\frac{1}{4}$.

Remark

Complex hyperbolic plane $H_{\mathbb{C}}^2$ is isometric to $G' = \text{Heis}^3 \rtimes_{\alpha} \mathbb{R}$ where $\alpha = \text{diag}(1, 1, 2)$ and Heis denotes the Heisenberg group. Therefore it is very close to G .

Theorem

$T^{2,p}(H_{\mathbb{C}}^2) = 0$ for $2 < p < 4$.

Proof of torsion comparison theorem

Use the gradient vectorfield ξ of a Busemann function and its flow ϕ_t , whose derivative is controlled by sectional curvature. For α a closed k -form in L^p ,

$$\phi_t^* \alpha = \alpha + d \left(\int_0^t \phi_s^* \iota_\xi \alpha \, ds \right)$$

has a limit as $t \rightarrow +\infty$ under the assumptions of the theorem. This boundary value map injects $H^{k,p}$ into a function space of closed forms on the ideal boundary, showing that $H^{k,p}$ is Hausdorff.

Proof of torsion comparison theorem

Use the gradient vectorfield ξ of a Busemann function and its flow ϕ_t , whose derivative is controlled by sectional curvature. For α a closed k -form in L^p ,

$$\phi_t^* \alpha = \alpha + d \left(\int_0^t \phi_s^* \iota_\xi \alpha \, ds \right)$$

has a limit as $t \rightarrow +\infty$ under the assumptions of the theorem. This boundary value map injects $H^{k,p}$ into a function space of closed forms on the ideal boundary, showing that $H^{k,p}$ is Hausdorff.

Proof of torsion vanishing for $H_{\mathbb{C}}^2$

For $p \notin \{4/3, 2, 4\}$, differential forms α on $H_{\mathbb{C}}^2$ split into components α_+ and α_- which are contracted (resp. expanded) by ϕ_t . Then

$$B_t : \alpha \mapsto \int_0^t \phi_s^* \iota_\xi \alpha_+ \, ds - \int_{-t}^0 \phi_s^* \iota_\xi \alpha_- \, ds$$

converges as $t \rightarrow +\infty$ to a bounded operator B on L^p . $P = 1 - dB - Bd$ retracts the L^p de Rham complex onto a complex of differential forms on Heis^3 with missing components and weakly regular coefficients. If $2 < p < 4$, this complex is nonzero in degrees 1 and 2, but it is so small that its cohomology can be shown to be Hausdorff.

Non-vanishing of torsion

Use Poincaré duality. Let $p' = p/p - 1$ denote the conjugate exponent. In order to prove that a closed k -form α is nonzero in cohomology, it suffices to construct a sequence ψ_j of $(n - k)$ -forms such that $\|d\psi_j\|_{L^{p'}}$ tends to zero but $\int \alpha \wedge \psi_j$ does not tend to zero.

Use Poincaré duality. Let $p' = p/p - 1$ denote the conjugate exponent. In order to prove that a closed k -form α is nonzero in cohomology, it suffices to construct a sequence ψ_j of $(n - k)$ -forms such that $\|d\psi_j\|_{L^{p'}}$ tends to zero but $\int \alpha \wedge \psi_j$ does not tend to zero.

In conclusion, we have used

- ▶ Poincaré duality.
- ▶ A deformation retraction of space onto a subspace, with controlled effect on the L^p -norms of forms. For certain ranges of p , this provides a boundary value.


Use Poincaré duality. Let $p' = p/p - 1$ denote the conjugate exponent. In order to prove that a closed k -form α is nonzero in cohomology, it suffices to construct a sequence ψ_j of $(n - k)$ -forms such that $\|d\psi_j\|_{L^{p'}}$ tends to zero but $\int \alpha \wedge \psi_j$ does not tend to zero.

In conclusion, we have used

- ▶ Poincaré duality.
- ▶ A deformation retraction of space onto a subspace, with controlled effect on the L^p -norms of forms. For certain ranges of p , this provides a boundary value.

Conjecture


- ▶ For rank 1 symmetric spaces, $T^{k,p} = 0$ except for at most 1 value of p in each degree.
- ▶ For higher rank symmetric spaces, $H^{k,p} = 0$ for $k < \text{rank}$, $T^{k,p} = 0$ for $k = \text{rank}$.
- ▶ For $k = \text{rank}$, $R^{k,p} \neq 0$ for p large, and $R^{k,p}$ is a function space on the maximal boundary.
- ▶ For each $p > 1$, there exists k such that $H^{k,p} \neq 0$.

L^p cohomology of $H_{\mathbb{C}}^2$ 

 non vanishing reduced cohomology

 non vanishing torsion

L^p -cohomology of $\mathbb{R}^{\times_{\text{diag}(1,1,2)}} \mathbb{R}^3$



 non vanishing reduced cohomology

 non vanishing torsion