

Lecture 10 Double integral

- Recall on the simple integral

We suppose given two reals $a < b$ and a function $f : [a, b] \rightarrow \mathbb{R}$. The integral $\int_a^b f(x) dx$ of f on the interval $[a, b]$, denoted also as $\int_{[a,b]} f$, is a real number satisfying the following properties :

- ★ Length. If $f(x) = 1$ for all x , then $\int_a^b dx = b - a$.
- ★ Linearity. If f and g are two functions $[a, b] \rightarrow \mathbb{R}$ and λ is a real number, we have $\int_a^b (f(x) + g(x)) dx = \int_a^b f(x) dx + \int_a^b g(x) dx$ and $\int_a^b (\lambda f(x)) dx = \lambda \int_a^b f(x) dx$.
- ★ Positivity. If f is a positive function, *id est* $f(x) \geq 0$ for all x , then the corresponding integral is positive : $\int_a^b f(x) dx \geq 0$. We remark that this property can be in defect if we do not suppose $a < b$.
- ★ Additivity relative to the domain (Chasles's relation). If $a < c < b$, then $\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$.

These properties can be extended to the double integral. It is not the case for the following ones.

- Specific properties of the simple integral

- ★ Fundamental theorem of Analysis and integration by parts. We suppose that f a continuous function $[a, b] \rightarrow \mathbb{R}$. For example, f can be a polynomial, a sinus or cosinus function, an exponential function, the absolute value or the composite of two continuous function. But f can not be the Heaviside function H proposed initially by Oliver Heaviside (1850-1925) and defined by $H(x) = 0$ for $x \leq 0$ and $H(x) = 1$ when $x > 0$. Then the mapping ψ defined according to $\psi(x) = \int_a^x f(\xi) d\xi$ is a derivable function of the variable x and $\frac{d}{dx} \left(\int_a^x f(\xi) d\xi \right) = f(x)$. In consequence, $\int_a^b \frac{df}{dx} d\xi = f(b) - f(a)$ if f is a continuously derivable function. The proof of this result introduces the function g defined according to $g(x) = f(x) - \int_a^x f(\xi) d\xi$. Then g is derivable and $\frac{dg}{dx} = 0$ on the interval $[a, b]$. In consequence, the function g is a constant function : $g(a) = g(b)$. This relation means that $f(a) = f(b) - \int_a^b f(\xi) d\xi$ and the result is proven. \square

In practice, this result is expressed as follows : If we suppose given a primitive function F of the function f (that is $\frac{dF}{dx} = f(x)$), then $\int_a^b f(x) dx = F(b) - F(a)$.

★ Change of variable. We suppose that the interval $[a, b]$ is parametrized by a one to one increasing function φ from $[\alpha, \beta]$ onto $[a, b] : x = \varphi(t)$ with $t \in [\alpha, \beta]$. Then

$$\int_a^b f(x) dx = \int_\alpha^\beta f(\varphi(t)) \varphi'(t) dt.$$

★ Calculus of surfaces. If the function f is positive from $[a, b]$ to \mathbb{R} (with $a < b$), then the integral $\int_a^b f(x) dx$ is equal to the area $|\Omega|$ of the domain Ω between the abscissae a and b on one hand, the abscissa axis and the curve $y = f(x)$ on the other hand :

$$\Omega = \{(x, y) \in \mathbb{R}^2, a \leq x \leq b, 0 \leq y \leq f(x)\}. \text{ Then we have } \int_a^b f(x) dx = |\Omega|.$$

• Fundamental properties of the double integral

We suppose given a bounded subset Ω of the plane \mathbb{R}^2 and a bounded function $f : \Omega \rightarrow \mathbb{R}$. The double integral of the function f on the domain Ω is a real number. It is noted $\int_\Omega f(x, y) dx dy$ or $\iint_\Omega f(x, y) dx dy$ and often more simply $\int_\Omega f dx dy$ or $\int_\Omega f$.

★ Surface : double integral of the function “one”. We introduce four real numbers a, b, c and d such that $a < b$ and $c < d$. We consider the rectangle $\Omega =]a, b[\times]c, d[$ of the plane \mathbb{R}^2 . The double integral of the function $f(x, y) \equiv 1$ is simply the area $(b - a)(d - c)$ of the rectangle : $\int_{]a, b[\times]c, d[} dx dy = (b - a)(d - c)$.

More generally, if Ω is a bounded part of the plane, that is if Ω is included in a large rectangle, the double integral on Ω of the function $f(x, y) \equiv 1$ is exactly the surface $|\Omega|$ of the domain : $\int_\Omega dx dy = |\Omega|$.

★ Linearity. We suppose that the double integral $\int_\Omega f(x, y) dx dy$ of the function f is known and we give a number λ . Then $\int_\Omega (\lambda f)(x, y) dx dy = \lambda \int_\Omega f(x, y) dx dy$. If we give also the double integral $\int_\Omega g(x, y) dx dy$ of the function g , then

$$\int_\Omega (f + g)(x, y) dx dy = \int_\Omega f(x, y) dx dy + \int_\Omega g(x, y) dx dy.$$

★ Positivity. We suppose that the function f is positive on $\Omega : f(x, y) \geq 0, \forall (x, y) \in \Omega$. Then $\int_\Omega f(x, y) dx dy \geq 0$. If $f \leq g$ on Ω that is $f(x, y) \leq g(x, y)$ for all $(x, y) \in \Omega$, then $\int_\Omega f(x, y) dx dy \leq \int_\Omega g(x, y) dx dy$ [exercice].

★ Additivity relative to the domain. We suppose the domain Ω is decomposed into a finite set of simpler sub-domains: $\Omega_i : \Omega = \cup_{i=1}^N \Omega_i$. Moreover, the intersection $\Omega_i \cap \Omega_j$ has a null area if $i \neq j : |\Omega_i \cap \Omega_j| = 0$. Then the integral on Ω of any function f is the sum of the integrals on each sub-domain Ω_i of this function : $\int_\Omega f(x, y) dx dy = \sum_{i=1}^N \int_{\Omega_i} f(x, y) dx dy$.

• Integral of a tiered function

We suppose given a decomposition of the domain Ω as above and a tiered function f on Ω , that is a constant function in each part $\Omega_i : \forall i = 1, \dots, N, \exists \lambda_i \in \mathbb{R}, \forall (x, y) \in \Omega_i, f(x, y) = \lambda_i$. The calculus of the integral of f on Ω is explicit : $\int_\Omega f(x, y) dx dy = \sum_{i=1}^N \lambda_i |\Omega_i|$ [exercice].

• Integral of a continuous function

We consider again a bounded domain $\Omega \subset \mathbb{R}^2$ and $f \in \mathcal{C}^0(\overline{\Omega})$ a continuous function on Ω up to the boundary. Then the integral of f on Ω is well defined ; it is a real number, or eventually a complex number.

• Fubini theorem [Guido Fubini (1879-1943)]

We consider a bounded domain $\Omega \subset \mathbb{R}^2$ and a bounded function defined on Ω with real or eventually complex values : $\exists M \geq 0, \forall (x, y) \in \Omega, |f(x, y)| \leq M$. Then the integral of the

absolute value of f is finite : $\iint_{\Omega} |f(x, y)| \, dx \, dy < \infty$. Moreover, the double integral of f in the domain Ω is well defined and we have the inequality $|\iint_{\Omega} f(x, y) \, dx \, dy| \leq \iint_{\Omega} |f(x, y)| \, dx \, dy$. The Fubini theorem says that it is always possible to integrate this function “in the order we want”.




Figure 1. Calculus of the double integral in the domain Ω , first by the integration of the function f relative to y between $\varphi_-(x)$ and $\varphi_+(x)$, and secondly by simple integration relative to x of the result, between a and b .

More precisely, if Ω is between two regular curves of the form $y = \varphi(x)$ as in Figure 1, that is $\Omega = \{(x, y) \in \mathbb{R}^2, a \leq x \leq b, \varphi_-(x) \leq y \leq \varphi_+(x)\}$, we have

$$\int_{\Omega} f(x, y) \, dx \, dy = \int_a^b dx \left[\int_{\varphi_-(x)}^{\varphi_+(x)} dy f(x, y) \right].$$

If Ω is included between two curves of the type $x = \psi(y)$ like in Figure 2, *id est* $\Omega = \{(x, y) \in \mathbb{R}^2, c \leq y \leq d, \psi_-(y) \leq x \leq \psi_+(y)\}$, we have $\int_{\Omega} f(x, y) \, dx \, dy = \int_c^d dy \left[\int_{\psi_-(y)}^{\psi_+(y)} dx f(x, y) \right]$.

When the domain Ω can be parametrized in two ways, the double integral can be computed by one relation or the other and we have

$$\int_{\Omega} f(x, y) \, dx \, dy = \int_a^b dx \left[\int_{\varphi_-(x)}^{\varphi_+(x)} dy f(x, y) \right] = \int_c^d dy \left[\int_{\psi_-(y)}^{\psi_+(y)} dx f(x, y) \right].$$

- A first example of using Fubini theorem

We suppose given two real numbers $a < b$ and an integrable function $g : [a, b] \rightarrow \mathbb{R}$. We suppose that the function g is positive : $\forall x \in [a, b], g(x) \geq 0$. We consider the domain $\Omega = \{(x, y) \in \mathbb{R}^2, a \leq x \leq b, 0 \leq y \leq g(x)\}$ considered previously with the study of simple integrals. Then the Fubini theorem, with $\varphi_- = 0$, $\varphi_+ = g$ and $f(x, y) \equiv 1$ for each $(x, y) \in \Omega$, allows to conclude that $|\Omega| = \int_a^b g(x) \, dx$. We recover the link between the simple integral and the calculus of surfaces.




Figure 2. Calculus of the double integral in the domain Ω , first by the integration of the function f relative to x between $\psi_-(y)$ and $\psi_+(y)$, and secondly by simple integration relative to y of the result, between c and d .

Exercices

- Rectangular domains

a) Let D be the domain $D = \{(x, y) \in \mathbb{R}^2, 0 \leq x \leq 1, 0 \leq y \leq 2\}$.

Compute the double integral $\int_D xy \, dx \, dy$. [1]

b) Same question with the domain $\Delta = \{(x, y) \in \mathbb{R}^2, 0 \leq x \leq \pi, 0 \leq y \leq \frac{\pi}{2}\}$ and the integral

$J = \int \int_{\Delta} x \sin(x+y) \, dx \, dy$. [$\pi - 2$]

- Double integral in a triangle

We suppose given two real numbers $a > 0$ and $b > 0$ and the triangle

$T = \{(x, y) \in \mathbb{R}^2, x \geq 0, y \geq 0, \frac{x}{a} + \frac{y}{b} \leq 1\}$. We consider the function $f(x, y) = x - y$.

a) Prove that the double integral of the absolute value $|f|$ on the triangle T is finite.

b) Compute the double integral $I = \iint_T f(x, y) \, dx \, dy$ with the first approach suggested by the Fubini theorem: $I = \iint_T f(x, y) \, dx \, dy = \int_0^a dx [\int_0^{b(1-x/a)} dy (x-y)]$ with an appropriate substitution of the question marks with an algebraic expression.

c) Compute the double integral I with the second approach suggested by the Fubini theorem: $I = \iint_T f(x, y) \, dx \, dy = \int_0^b dy [\int_0^{a(1-y/b)} dx (x-y)]$. The question marks will be replaced with a correct algebraic expression.

d) Compare your results of questions b) and c). Are they identical? [$\frac{1}{6} ab(a-b)$]

APPLIED MATHEMATICS

- Exchanging the order of integration

We consider a function f well defined for all reals x and y .

Complete the two expressions of the double integral $\int_0^1 dy \int_y^{\sqrt{y}} dx f(x, y) = \int_?^? dx \int_?^? dy f(x, y)$ obtained after the exchange of the order of the integrals.

[remark that $\Omega = \{(x, y) \in \mathbb{R}^2, 0 \leq y \leq 1, y \leq x \leq \sqrt{y}\}$]

- Inequalities

a) If $f \leq g$ on Ω that is $f(x, y) \leq g(x, y)$ for all $(x, y) \in \Omega$, prove that we have the following inequality between numbers $\int_{\Omega} f(x, y) dx dy \leq \int_{\Omega} g(x, y) dx dy$.

If f is a bounded function in the bounded domain Ω , we introduce the two positive functions $f^+ = \max(f, 0)$ and $f^- = \max(-f, 0)$.

b) Prove that $f = f^+ - f^-$ and $|f| = f^+ + f^-$.

c) Deduce from the previous question that we have $|\iint_{\Omega} f(x, y) dx dy| \leq \iint_{\Omega} |f(x, y)| dx dy$.