

Infrared and ultraviolet problem for the Nelson model with variable coefficients

C. Gérard¹, F. Hiroshima², A. Panati³, and A. Suzuki²

¹Département de Mathématiques, Université de Paris XI, 91405 Orsay
Cedex France

²Department of Mathematics, University of Kyushu, 6-10-1, Hakozaki,
Fukuoka, 812-8581, Japan

³PHYMAT Université Toulon-Var 83957 La Garde Cedex France

April 2010

Abstract

We consider the Nelson model with variable coefficients and investigate the problem of existence of a ground state and the removal of the ultraviolet cutoff. Nelson models with variable coefficients arise when one replaces in the usual Nelson model the flat Minkowski metric by a static metric, allowing also the boson mass to depend on position. A physical example is obtained by quantizing the Klein-Gordon equation on a static space-time coupled with a non-relativistic particle. We investigate the existence of a ground state of the Hamiltonian in the presence of the infrared problem, i.e. assuming that the boson mass tends to 0 at infinity. We also study the removal of the ultraviolet cutoff, which allows to construct a model with a local interaction.

1 Introduction

In this paper we consider a class of quantum field theory Hamiltonians that we call *variable coefficients Nelson models*. These models are natural extensions of the usual Nelson model to the case when the Minkowski metric is replaced by a general static metric and the boson mass is position dependent. In this introduction we describe these models and summarize the results of this paper.

AMS 2010 *Subject Classification*. 81T10, 81T20, 81Q10, 58C40

Key words and phrases. Nelson model, static space-times, ground state, ultraviolet limit

1.1 The Nelson model on Minkowski space-time

The *Nelson* model describes a scalar bosonic field linearly coupled to a quantum mechanical particle. It is formally defined by the Hamiltonian

$$H = \frac{1}{2}p^2 + W(q) + \frac{1}{2} \int_{\mathbb{R}^3} \pi^2(\mathbf{x}) + (\nabla\varphi(\mathbf{x}))^2 + m^2\varphi^2(\mathbf{x})d\mathbf{x} + \int_{\mathbb{R}^3} \varphi(\mathbf{x})\rho(\mathbf{x} - q)d\mathbf{x},$$

where ρ denotes a cutoff function, p , q denote the position and momentum of the particle, $W(q)$ is an external potential and $\varphi(\mathbf{x})$, $\pi(\mathbf{x})$ are the canonical field position and momentum.

The Nelson model arises from the quantization of the following coupled Klein-Gordon and Newton system:

$$(1.1) \quad \begin{cases} (\square + m^2)\varphi(t, \mathbf{x}) = -\rho(\mathbf{x} - q_t), \\ \ddot{q}_t = -\nabla_q W(q_t) - \int \varphi(t, \mathbf{x}) \nabla_x \rho(\mathbf{x} - q_t) d\mathbf{x}, \end{cases}$$

where \square denotes the d'Alembertian on the Minkowski space-time \mathbb{R}^{1+3} . The cutoff function ρ plays the role of an ultraviolet cutoff and amounts to replacing the quantum mechanical point particle by a charge density.

To distinguish the Nelson model on Minkowski space-time from its generalizations that will be described later in the introduction, we will call it the *usual* (or *constant coefficients*) *Nelson model*.

The initial interest of the Nelson model [Ne] was that it is the simplest non trivial QFT model for which the ultraviolet limit, (amounting to replace the cutoff function ρ by the delta function δ), can be performed by relatively easy arguments. With the ultraviolet cutoff removed, the Nelson model becomes a local QFT model.

Even with an ultraviolet cutoff, the rigorous study of the Nelson model is of much interest, and quite a lot of efforts were devoted to the rigorous analysis of several of its aspects (see [AHH], [BFS], [BHLMS], [H], [LMS], [Sp]).

One of them, which will also be our main interest in this paper, is the question of the *existence of a ground state*. Obviously the fact that H has a ground state is an important physical property of the Nelson model. For example a consequence of the existence of a ground state is that *scattering states* can quite easily be constructed. These states describe the ground state of H with a finite number of additional asymptotically free bosons.

When H has no ground state one usually speaks of the *infrared problem* or *infrared divergence*. The infrared problem arises when the emission probability of bosons becomes infinite with increasing wave length. If the infrared problem occurs, the scattering theory has to be modified: all scattering states contain an infinite number of low energy (soft) bosons (see eg [DG3]).

For the usual Nelson model the answer to this question is well known: one assumes a stability condition (see Subsect. 4.5), implying that states with energy close to the bottom of the spectrum are localized in the particle position. Then if the bosons are massive i.e. if $m > 0$ H has a ground state (see eg [G]). On the contrary if $m = 0$ and $\int \rho(x)dx \neq 0$ then H has no ground state (see [DG3]).

1.2 The Nelson model with variable coefficients

In this paper we will study generalizations of the usual Nelson model, obtained by replacing the free Laplacian $-\Delta_x$ by a general second order differential operator and the constant mass term m by a function $m(x)$. We set:

$$h := - \sum_{1 \leq j, k \leq d} c(x)^{-1} \partial_j a^{jk}(x) \partial_k c(x)^{-1} + m^2(x),$$

for a Riemannian metric $a_{jk} dx^j dx^k$ and two functions $c(x)$, $m(x) > 0$, and consider the generalization of (1.1):

$$(1.2) \quad \begin{cases} \partial_t^2 \phi(t, x) + h\phi(t, x) + \rho(x - q_t) = 0, \\ \ddot{q}_t = -\nabla_x W(q_t) - \int_{\mathbb{R}^3} \phi(t, x) \nabla_x \rho(x - q_t) |g|^{\frac{1}{2}} d^3x. \end{cases}$$

Quantizing the field equations (1.2), we obtain a Hamiltonian H acting on the Hilbert space $L^2(\mathbb{R}^3) \otimes \Gamma_s(L^2(\mathbb{R}^3))$ (see Sect. 3), which we call a *Nelson Hamiltonian with variable coefficients*. Formally H is defined by the following expression:

$$(1.3) \quad \begin{aligned} H = & \frac{1}{2} p^2 + W(q) \\ & + \frac{1}{2} \int_{\mathbb{R}^3} \pi^2(x) + \sum_{jk} \partial_j c(x)^{-1} \varphi(x) a^{jk}(x) \partial_k c(x)^{-1} \varphi(x) + m^2(x) \varphi^2(x) dx \\ & + \int_{\mathbb{R}^3} \varphi(x) \rho(x - q) dx. \end{aligned}$$

The main example of a variable coefficients Nelson model is obtained by replacing in the usual Nelson model the flat Minkowski metric on \mathbb{R}^{1+3} by a *static* Lorentzian metric, and by allowing also the mass m to be position dependent. Recall that a static metric on \mathbb{R}^{1+3} is of the form

$$g_{\mu\nu}(x) dx^\mu dx^\nu = -\lambda(x) dt dt + \lambda(x)^{-1} h_{\alpha\beta}(x) dx^\alpha dx^\beta,$$

where $x = (t, x) \in \mathbb{R}^{1+3}$, $\lambda(x) > 0$ is a smooth function, and $h_{\alpha,\beta}(x)$ is a Riemannian metric on \mathbb{R}^3 . We show in Subsect. 2.3 that the natural Lagrangian for a point particle coupled to a scalar field on (\mathbb{R}^{1+3}, g) leads (after a change of field variables) to the system (1.2).

1.3 The infrared problem

Assuming reasonable hypotheses on the matrix $[a^{jk}](x)$ and the functions $c(x)$, $m(x)$ it is easy to see that the formal expression (1.3) can be rigorously defined as a bounded below selfadjoint operator H .

The first question we address in this paper is the problem of existence of a ground state for H . Variable coefficients Nelson models are examples of an abstract class of QFT Hamiltonians called *abstract Pauli-Fierz Hamiltonians* (see eg [G], [BD] and Subsect. 4.1). If ω is the *one-particle energy*, the constant $m := \inf \sigma(\omega)$ can be called the (rest) mass of the bosonic field, and abstract Pauli-Fierz Hamiltonians fall naturally into two classes: massive models if $m > 0$ and massless if $m = 0$.

For massive models, H typically has a ground state, if we assume either that the quantum particle is confined or a stability condition (see Subsect. 4.5). In this paper we concentrate on the massless case and hence our typical assumption will be that

$$\lim_{x \rightarrow \infty} m(x) = 0.$$

It follows that bosons of arbitrarily small energy may be present. The main result of this paper is that the existence or non-existence of a ground state for H depends on the rate of decay of the function $m(x)$. In fact we show in Thm. 4.1 that if

$$m(x) \geq a \langle x \rangle^{-1}, \text{ for some } a > 0,$$

and if the quantum particle is confined, then H has a ground state. In a subsequent paper [GHPS2], we will show that if

$$0 \leq m(x) \leq C \langle x \rangle^{-1-\epsilon}, \text{ for some } \epsilon > 0,$$

then H has no ground state. Therefore Thm. 4.1 is sharp with respect to the decay rate of the mass at infinity.

(If $h = -\Delta + \lambda m^2(x)$ for $m(x) \in O(\langle x \rangle^{-3/2})$ and the coupling constant λ is sufficiently small the same result is shown in [GHPS1]).

1.4 Removal of the UV cutoff

As explained in Subsect. 1.1 the ultraviolet limit of the constant coefficients Nelson model was rigorously constructed long ago by Nelson. We consider also the same question for variable coefficients Nelson models. Denoting by H^κ the Nelson Hamiltonian H for the cutoff function $\rho_\kappa(x) = \kappa^3 \rho(\kappa x)$, we construct a particle potential $E^\kappa(q)$ such that $H^\kappa - E^\kappa(q)$ converge in strong resolvent sense to a bounded below selfadjoint operator H^∞ (see Thm. 5.5).

The removal of the UV cutoff involves as in the constant coefficients case a sequence of unitary dressing operators U^κ . In contrary to the constant coefficients case, where all computations can be conveniently done in momentum space (after conjugation by Fourier transform), we have to use instead *pseudodifferential calculus*. Some of the rather advanced facts on pseudodifferential calculus which we will need are recalled in Appendix B.

1.5 Notation

We collect here some notation for the reader's convenience.

If $x \in \mathbb{R}^d$, we set $\langle x \rangle = (1 + x^2)^{\frac{1}{2}}$.

The domain of a linear operator A on some Hilbert space \mathcal{H} will be denoted by $\text{Dom}A$, and its spectrum by $\sigma(A)$.

If \mathfrak{h} is a Hilbert space, the *bosonic Fock space* over \mathfrak{h} denoted by $\Gamma_s(\mathfrak{h})$ is

$$\Gamma_s(\mathfrak{h}) := \bigoplus_{n=0}^{\infty} \otimes_s^n \mathfrak{h}.$$

We denote by $a^*(h)$, $a(h)$ for $h \in \mathfrak{h}$ the *creation/annihilation operators* acting on $\Gamma_s(\mathfrak{h})$. The (Segal) *field operators* $\phi(h)$ are defined as $\phi(h) := \frac{1}{\sqrt{2}}(a^*(h) + a(h))$.

If \mathcal{K} is another Hilbert space and $v \in B(\mathcal{K}, \mathcal{K} \otimes \mathfrak{h})$, then one defines the operators $a^*(v)$, $a(v)$ as unbounded operators on $\mathcal{K} \otimes \Gamma_s(\mathfrak{h})$ by:

$$\begin{aligned} a^*(v) \Big|_{\mathcal{K} \otimes \otimes_s^n \mathfrak{h}} &:= \sqrt{n+1} \left(\mathbb{1}_{\mathcal{K}} \otimes \mathcal{S}_{n+1} \right) \left(v \otimes \mathbb{1}_{\otimes_s^n \mathfrak{h}} \right), \\ a(v) &:= \left(a^*(v) \right)^*, \\ \phi(v) &:= \frac{1}{\sqrt{2}}(a(v) + a^*(v)). \end{aligned}$$

They satisfy the estimates

$$(1.4) \quad \|a^\sharp(v)(N+1)^{-\frac{1}{2}}\| \leq \|v\|,$$

where $\|v\|$ is the norm of v in $B(\mathcal{K}, \mathcal{K} \otimes \mathfrak{h})$.

If b is a selfadjoint operator on \mathfrak{h} its second quantization $d\Gamma(b)$ is defined as:

$$d\Gamma(b) \Big|_{\otimes_s^n \mathfrak{h}} := \sum_{j=1}^n \underbrace{\mathbb{1} \otimes \cdots \otimes \mathbb{1}}_{j-1} \otimes b \otimes \underbrace{\mathbb{1} \otimes \cdots \otimes \mathbb{1}}_{n-j}.$$

2 The Nelson model on static space-times

In this section we discuss the Nelson model on static space-times, which is the main example of Hamiltonians that will be studied in the rest of the paper. It is convenient to start with the Lagrangian framework.

2.1 Klein-Gordon equation on static space-times

Let $g_{\mu\nu}(x)$ be a Lorentzian metric of signature $(-, +, +, +)$ on \mathbb{R}^{1+3} . Set $|g| = \det[g_{\mu\nu}]$, $[g^{\mu\nu}] = [g_{\mu\nu}]^{-1}$. Consider the Lagrangian

$$L_{\text{free}}(\phi)(x) = \frac{1}{2} \partial_\mu \phi(x) g^{\mu\nu}(x) \partial_\nu \phi(x) + \frac{1}{2} m^2(x) \phi^2(x),$$

for a function $m : \mathbb{R}^4 \rightarrow \mathbb{R}^+$ and the associated action:

$$S_{\text{field}}(\phi) = \int_{\mathbb{R}^4} L_{\text{free}}(\phi)(x) |g|^{\frac{1}{2}}(x) d^4x,$$

where $\phi : \mathbb{R}^4 \rightarrow \mathbb{R}$. The Euler-Lagrange equations yield the *Klein-Gordon equation*:

$$\square_g \phi + m^2(x)\phi = 0,$$

for

$$\square_g = -|g|^{-\frac{1}{2}} \partial_\mu |g|^{\frac{1}{2}} g^{\mu\nu} \partial_\nu.$$

Usually one has

$$\frac{1}{2}m^2(x) = \frac{1}{2}(m^2 + \theta R(x)),$$

where $m \geq 0$ is the mass and $R(x)$ is the scalar curvature of the metric $g_{\mu\nu}$, (assuming of course that the function on the right is positive). In particular if $m = 0$ and $\theta = \frac{1}{6}$ one obtains the so-called conformal wave equation.

We set $x = (t, \mathbf{x}) \in \mathbb{R}^{1+3}$. The metric $g_{\mu\nu}$ is *static* if:

$$g_{\mu\nu}(x)dx^\mu dx^\nu = -\lambda(x)dtdt + \lambda(x)^{-1}h_{\alpha\beta}(x)dx^\alpha dx^\beta,$$

where $\lambda(x) > 0$ is a smooth function and $h_{\alpha\beta}$ is a Riemannian metric on \mathbb{R}^3 . We assume also that $m^2(x) = m^2(\mathbf{x})$ is independent on t .

Setting $\phi(t, \mathbf{x}) = \lambda|h|^{-1/4}\tilde{\phi}(t, \mathbf{x})$, we obtain that $\tilde{\phi}(t, \mathbf{x})$ satisfies the equation:

$$\partial_t^2 \tilde{\phi} - \lambda|h|^{-1/4} \partial_\alpha |h|^{\frac{1}{2}} h^{\alpha\beta} \partial_\beta |h|^{-1/4} \lambda \tilde{\phi} + m^2 \lambda \tilde{\phi} = 0.$$

We note that $|h|^{-1/4} \partial_\alpha |h|^{\frac{1}{2}} h^{\alpha\beta} \partial_\beta |h|^{-1/4}$ is (formally) self-adjoint on $L^2(\mathbb{R}^3, dx)$ and is the Laplace-Beltrami operator Δ_h associated to the Riemannian metric $h_{\alpha\beta}$ (after the usual density change $u \mapsto |h|^{1/4}u$ to work on the Hilbert space $L^2(\mathbb{R}^3, dx)$).

2.2 Klein-Gordon field coupled to a non-relativistic particle

We now couple the Klein-Gordon field to a non-relativistic particle. We fix a mass $M > 0$, a charge density $\rho : \mathbb{R}^3 \rightarrow \mathbb{R}^+$ with $q = \int_{\mathbb{R}^3} \rho(y) d^3y \neq 0$ and a real potential $W : \mathbb{R}^3 \rightarrow \mathbb{R}$. The action for the coupled system is

$$S = S_{\text{part}} + S_{\text{field}} + S_{\text{int}},$$

for

$$\begin{aligned} S_{\text{part}} &= \int_{\mathbb{R}} \frac{M}{2} |\dot{\mathbf{x}}(t)|^2 - W(\mathbf{x}(t)) dt, \\ S_{\text{int}} &= \int_{\mathbb{R}^4} \phi(t, \mathbf{x}) \rho(\mathbf{x} - \mathbf{x}(t)) |g|^{\frac{1}{2}}(x) d^4x. \end{aligned}$$

The Euler-Lagrange equations are:

$$\begin{cases} \square_g \phi(t, \mathbf{x}) + m^2(t, \mathbf{x})\phi(t, \mathbf{x}) + \rho(\mathbf{x} - \mathbf{x}(t)) = 0, \\ M\ddot{\mathbf{x}}(t) = -\nabla_{\mathbf{x}} W(\mathbf{x}(t)) - \int_{\mathbb{R}^3} \phi(t, \mathbf{x}) \nabla_{\mathbf{x}} \rho(\mathbf{x} - \mathbf{x}(t)) |g|^{\frac{1}{2}} d^3\mathbf{x}. \end{cases}$$

Doing the same change of field variables as in Subsect. 2.1 and deleting the tildes, we obtain the system:

$$(2.1) \quad \begin{cases} \partial_t^2 \phi - \lambda \Delta_h \lambda \phi + m^2 \lambda \phi + \rho(\mathbf{x} - \mathbf{x}(t)) = 0, \\ M\ddot{\mathbf{x}}(t) = -\nabla W(\mathbf{x}(t)) - \int_{\mathbb{R}^3} \phi(t, \mathbf{x}) \nabla \rho(\mathbf{x} - \mathbf{x}(t)) d^3\mathbf{x}. \end{cases}$$

2.3 The Nelson model on a static space-time

If the metric is static, the equations (2.1) are clearly Hamiltonian equations for the classical Hamiltonian $H = H_{\text{part}} + H_{\text{field}} + H_{\text{int}}$, where:

$$\begin{aligned} H_{\text{part}}(\mathbf{x}, \xi) &= \frac{1}{2M}\xi^2 + W(\mathbf{x}), \\ H_{\text{field}}(\varphi, \pi) &= \frac{1}{2} \int_{\mathbb{R}^3} \pi^2(\mathbf{x}) - \varphi(\mathbf{x})\lambda(\mathbf{x})\Delta_h\lambda(\mathbf{x})\varphi(\mathbf{x}) + m^2(\mathbf{x})\lambda(\mathbf{x})\varphi^2(\mathbf{x})d\mathbf{x}, \\ H_{\text{int}}(\mathbf{x}, \xi, \varphi, \pi) &= \int_{\mathbb{R}^3} \rho(\mathbf{y} - \mathbf{x})\varphi(\mathbf{y})d\mathbf{y}. \end{aligned}$$

The classical phase space is as usual $\mathbb{R}^3 \times \mathbb{R}^3 \times L^2_{\mathbb{R}}(\mathbb{R}^3) \times L^2_{\mathbb{R}}(\mathbb{R}^3)$, with the symplectic form

$$(\mathbf{x}, \xi, \varphi, \pi)\omega(\mathbf{x}', \xi', \varphi', \pi') = \mathbf{x} \cdot \xi' - \mathbf{x}' \cdot \xi + \int_{\mathbb{R}^3} \varphi(\mathbf{x})\pi'(\mathbf{x}) - \pi(\mathbf{x})\varphi'(\mathbf{x})d\mathbf{x}.$$

The usual quantization scheme leads to the Hilbert space:

$$L^2(\mathbb{R}^3, d\mathbf{y}) \otimes \Gamma_s(L^2(\mathbb{R}^3, d\mathbf{x})),$$

where $\Gamma_s(\mathfrak{h})$ is the bosonic Fock space over the one-particle space \mathfrak{h} , and to the quantum Hamiltonian:

$$H = \left(-\frac{1}{2}\Delta_{\mathbf{y}} + W(\mathbf{y})\right) \otimes \mathbb{1} + \mathbb{1} \otimes d\Gamma(\omega) + \frac{1}{\sqrt{2}} \left(a^*(\omega^{-\frac{1}{2}}\rho(\cdot - \mathbf{y})) + a(\omega^{-\frac{1}{2}}\rho(\cdot - \mathbf{y})) \right),$$

where

$$\omega = (-\lambda\Delta_h\lambda + m^2\lambda)^{\frac{1}{2}},$$

$d\Gamma(\omega)$ is the usual second quantization of ω and $a^*(f)$, $a(f)$ are the creation/annihilation operators on $\Gamma_s(L^2(\mathbb{R}^3, d\mathbf{x}))$.

3 The Nelson Hamiltonian with variable coefficients

In this section we define the Nelson model with variable coefficients that will be studied in the rest of the paper. We will deviate slightly from the notation in Sect. 2 by denoting by $x \in \mathbb{R}^3$ (resp. $X \in \mathbb{R}^3$) the boson (resp. electron) position. As usual we set $D_x = i^{-1}\nabla_x$, $D_X = i^{-1}\nabla_X$.

3.1 Electron Hamiltonian

We define the electron Hamiltonian as:

$$K := K_0 + W(X),$$

where

$$K_0 = \sum_{1 \leq j, k \leq 3} D_{X_j} A^{jk}(X) D_{X_k},$$

acting on $\mathcal{K} := L^2(\mathbb{R}^3, dX)$, where:

$$(E1) \quad C_0 \mathbb{1} \leq [A^{jk}(X)] \leq C_1 \mathbb{1}, \quad C_0 > 0.$$

We assume that $W(X)$ is a real potential such that $K_0 + W$ is essentially selfadjoint and bounded below. We denote by K the closure of $K_0 + W$. Later we will assume the following *confinement condition* :

$$(E2) \quad W(X) \geq C_0 \langle X \rangle^{2\delta} - C_1, \quad \text{for some } \delta > 0.$$

Physically this condition means that the electron is confined. As is well known (see eg [GLL]) for the question of existence of a ground state, this condition can be replaced by a *stability condition*, meaning that states near the bottom of the spectrum of the Hamiltonian are confined in the electronic variables by energy conservation.

We will discuss the extension of our results when one assume the stability condition in Subsect. 4.5.

3.2 Field Hamiltonian

Let:

$$\begin{aligned} h_0 &:= - \sum_{1 \leq j, k \leq d} c(x)^{-1} \partial_j a^{jk}(x) \partial_k c(x)^{-1}, \\ h &:= h_0 + m^2(x), \end{aligned}$$

with a^{jk} , c , m are real functions and:

$$\begin{aligned} (B1) \quad & C_0 \mathbb{1} \leq [a^{jk}(x)] \leq C_1 \mathbb{1}, \quad C_0 \leq c(x) \leq C_1, \quad C_0 > 0, \\ & \partial_x^\alpha a^{jk}(x) \in O(\langle x \rangle^{-1}), \quad |\alpha| \leq 1, \quad \partial_x^\alpha c(x) \in O(1), \quad |\alpha| \leq 2, \\ & \partial_x^\alpha m(x) \in O(1), \quad |\alpha| \leq 1. \end{aligned}$$

Clearly h is selfadjoint on $H^2(\mathbb{R}^3)$ and $h \geq 0$. The *one-particle space* and *one-particle energy* are:

$$\mathfrak{h} := L^2(\mathbb{R}^3, dx), \quad \omega := h^{\frac{1}{2}}.$$

The constant:

$$\inf \sigma(\omega) =: m \geq 0,$$

can be viewed as the *mass* of the scalar bosons.

The following lemma is easy;

Lemma 3.1 (1) *One has $\text{Ker} \omega = \{0\}$,*

(2) *Assume in addition to (B1) that $\lim_{x \rightarrow \infty} m(x) = 0$. Then $\inf \sigma(\omega) = 0$.*

Proof. It follows from (B1) that

$$(u|hu) \leq C_1(c^{-1}u| -\Delta c^{-1}u) + (c^{-1}u|c^{-1}m^2u), \quad u \in H^2(\mathbb{R}^3).$$

Therefore if $hu = 0$ u is constant. It follows also from (B1) that $c(x)^{-1}$ preserves $H^2(\mathbb{R}^3)$. Therefore by the variational principle

$$m^2 = \inf \sigma(h) \leq C_1 \inf \sigma(-\Delta + c^{-2}(x)m^2(x)) = 0.$$

This proves (2). \square

The Nelson Hamiltonian defined below will be called *massive* (resp. *massless*) if $m > 0$ (resp. $m = 0$.) The field Hamiltonian is

$$d\Gamma(\omega),$$

acting on the bosonic Fock space $\Gamma_s(\mathfrak{h})$.

3.3 Nelson Hamiltonian

Let $\rho \in S(\mathbb{R}^3)$, with $\rho \geq 0$, $q = \int_{\mathbb{R}^3} \rho(y)dy \neq 0$. We set:

$$\rho_X(x) = \rho(x - X)$$

and define the *UV cutoff fields* as:

$$(3.1) \quad \varphi_\rho(X) := \phi(\omega^{-\frac{1}{2}}\rho_X),$$

where for $f \in \mathfrak{h}$, $\phi(f)$ is the Segal field operator:

$$\phi(f) := \frac{1}{\sqrt{2}} (a^*(f) + a(f)).$$

Note that setting

$$\varphi(X) := \phi(\omega^{-\frac{1}{2}}\delta_X),$$

one has $\varphi_\rho(X) = \int \varphi(X - Y)\rho(Y)dY$.

Remark 3.2 *One can think of another definition of UV cutoff fields, namely:*

$$\tilde{\varphi}_\chi(X) := \phi(\omega^{-\frac{1}{2}}\chi(\omega)\delta_X),$$

for $\chi \in S(\mathbb{R})$, $\chi(0) = 1$. In the constant coefficients case where $h = -\Delta$ both definitions are equivalent. In the variable coefficients case the natural definition (3.1) is much more convenient.

The *Nelson Hamiltonian* is:

$$(3.2) \quad H := K \otimes \mathbb{1} + \mathbb{1} \otimes d\Gamma(\omega) + \varphi_\rho(X),$$

acting on

$$\mathcal{H} = \mathcal{K} \otimes \Gamma_s(\mathfrak{h}).$$

Set also:

$$H_0 := K \otimes \mathbb{1} + \mathbb{1} \otimes d\Gamma(\omega),$$

which is selfadjoint on its natural domain. The following lemma is standard.

Lemma 3.3 *Assume hypotheses (E1), (B1). Then H is selfadjoint and bounded below on $D(H_0)$.*

Proof. it suffices to apply results on abstract Pauli-Fierz Hamiltonians (see eg [GGM, Sect.4]). H is an abstract Pauli-Fierz Hamiltonian with coupling operator $v \in B(\mathcal{K}, \mathcal{K} \otimes \mathfrak{h})$ equal to:

$$L^2(\mathbb{R}^3, dX) \ni u \mapsto \omega^{-\frac{1}{2}} \rho(x - X)u(X) \in L^2(\mathbb{R}^3, dX) \otimes L^2(\mathbb{R}^3, dx)$$

Applying [GGM, Corr. 4.4], it suffices to check that $\omega^{-\frac{1}{2}}v \in B(\mathcal{K}, \mathcal{K} \otimes \mathfrak{h})$. Now

$$\|\omega^{-\frac{1}{2}}v\|_{B(\mathcal{K}, \mathcal{K} \otimes \mathfrak{h})} = \left(\sup_{X \in \mathbb{R}^3} \|\omega^{-1} \rho_X\|^2 \right)^{\frac{1}{2}}$$

Using that $h \geq CD_x^2$ and the Kato-Heinz inequality, we obtain that $\omega^{-2} \leq C|D_x|^{-2}$, hence it suffices to check that the map

$$L^2(\mathbb{R}^3, dX) \ni u \mapsto |D_x|^{-1} \rho(x - X)u(X) \in L^2(\mathbb{R}^3, dX) \otimes L^2(\mathbb{R}^3, dx)$$

is bounded, which is well known. \square

4 Existence of a ground state

In this section we will prove our main result about the existence of a ground state for variable coefficients Nelson Hamiltonians. This result will be deduced from an abstract existence result extending the one in [BD], whose proof is outlined in Subsects. 4.1, 4.2 and 4.3.

Theorem 4.1 *Assume hypotheses (E1), (B1). Assume in addition that:*

$$m(x) \geq a\langle x \rangle^{-1}, \text{ for some } a > 0,$$

and (E2) for some $\delta > \frac{3}{2}$. Then $\inf \sigma(H)$ is an eigenvalue.

Remark 4.2 *The condition $\delta > \frac{3}{2}$ in Thm. 4.1 comes from the operator bound $\omega^{-3} \leq C\langle x \rangle^{3+\epsilon}$, $\forall \epsilon > 0$ proved in Thm. A.8.*

Remark 4.3 From Lemma 3.1 we know that $\inf \sigma(\omega) = 0$ if $\lim_{x \rightarrow \infty} m(x) = 0$. Therefore the Nelson Hamiltonian can be massless using the terminology of Subsect. 3.2.

Remark 4.4 In a subsequent paper [GHPS2] we will show that if

$$0 \leq m(x) \leq C \langle x \rangle^{-1-\epsilon}, \text{ for some } \epsilon > 0,$$

then H has no ground state. Therefore the result of Thm. 4.1 is sharp with respect to the decay rate of the mass at infinity.

4.1 Abstract Pauli-Fierz Hamiltonians

In [BD], Bruneau and Dereziński study the spectral theory of abstract Pauli-Fierz Hamiltonians of the form

$$H = K \otimes \mathbb{1} + \mathbb{1} \otimes d\Gamma(\omega) + \phi(v),$$

acting on the Hilbert space $\mathcal{H} = \mathcal{K} \otimes \Gamma_s(\mathfrak{h})$, where \mathcal{K} is the Hilbert space for the small system and \mathfrak{h} the one-particle space for the bosonic field. The Hamiltonian H is called massive (resp. massless) if $\inf \sigma(\omega) > 0$ (resp. $\inf \sigma(\omega) = 0$). Among other results they prove the existence of a ground state for H if v is infrared regular.

Although most of their hypotheses are natural and essentially optimal, we cannot directly apply their abstract results to our situation. In fact they assume (see [BD, Assumption E]) that the one-particle space \mathfrak{h} equals $L^2(\mathbb{R}^d, dk)$ and the one-particle energy ω is the multiplication operator by a function $\omega(k)$ which is positive, with $\nabla\omega$ bounded, and $\lim_{k \rightarrow \infty} \omega(k) = +\infty$. This assumption on the one-particle energy is only needed to prove an HVZ theorem for massive (or massless with an infrared cutoff) Pauli-Fierz Hamiltonians.

In our case this assumption could be deduced (modulo unitary equivalence) from the spectral theory of h . For example it would suffice to know that h is unitarily equivalent to $-\Delta$. This last property would follow from the absence of eigenvalues for h and from the scattering theory for the pair $(h, -\Delta)$ and require additional decay properties of the $[a^{ij}](x)$, $m(x)$ and of some of their derivatives.

We will replace it by more geometric assumptions on ω (see hypothesis (4.4) below), similar to those introduced in [GP], where abstract bosonic QFT Hamiltonians were considered. Since we do not aim for generality, our hypotheses on the coupling operator v are stronger than necessary, but lead to simpler proofs. Also most of the proofs will be only sketched.

Let $\mathfrak{h}, \mathcal{K}$ two Hilbert spaces and set $\mathcal{H} = \mathcal{K} \otimes \Gamma_s(\mathfrak{h})$.

We fix selfadjoint operators $K \geq 0$ on \mathcal{K} and $\omega \geq 0$ on \mathfrak{h} . We set

$$\inf \sigma(\omega) =: m \geq 0.$$

If $m = 0$ one has to assume additionally that $\text{Ker}\omega = \{0\}$ (see Remark 4.5 for some explanation of this fact).

Remark 4.5 *It \mathcal{X} is a real Hilbert space and ω is a selfadjoint operator on \mathcal{X} , the condition $\text{Ker}\omega = \{0\}$ is well known to be necessary to have a stable quantization of the abstract Klein-Gordon equation $\partial_t^2\phi(t) + \omega^2\phi(t) = 0$ where $\phi(t) : \mathbb{R} \rightarrow \mathcal{X}$.*

If $\text{Ker}\omega \neq \{0\}$ the phase space $\mathcal{Y} = \mathcal{X} \oplus \mathcal{X}$ for the Klein-Gordon equation splits into the symplectic direct sum $\mathcal{Y}_{\text{reg}} \oplus \mathcal{Y}_{\text{sing}}$, for $\mathcal{Y}_{\text{reg}} = \text{Ker}\omega^\perp \oplus \text{Ker}\omega^\perp$, $\mathcal{Y}_{\text{sing}} = \text{Ker}\omega \oplus \text{Ker}\omega$, both symplectic spaces being invariant under the symplectic evolution associated to the Klein-Gordon equation. On \mathcal{Y}_{reg} one can perform the stable quantization. On $\mathcal{Y}_{\text{sing}}$, if for example $\text{Ker}\omega$ is d -dimensional, the quantization leads to the Hamiltonian $-\Delta$ on $L^2(\mathbb{R}^d)$. Clearly any perturbation of the form $\phi(f)$ for $\mathbb{1}_{\{0\}}(\omega)f \neq 0$ will make the Hamiltonian unbounded from below.

So we will always assume that

$$(4.1) \quad \omega \geq 0, \text{Ker}\omega = \{0\}.$$

Let $H_0 = \mathcal{K} \otimes \mathbb{1} + \mathbb{1} \otimes d\Gamma(\omega)$. We fix also a coupling operator v such that:

$$(4.2) \quad v \in B(\mathcal{K}, \mathcal{K} \otimes \mathfrak{h}).$$

The quadratic form $\phi(v) = a(v) + a^*(v)$ is well defined for example on $\mathcal{K} \otimes \text{Dom}N^{\frac{1}{2}}$. We will also assume that:

$$(4.3) \quad \omega^{-\frac{1}{2}}v(K+1)^{-\frac{1}{2}} \text{ is compact.}$$

Proposition 4.6 ([BD] Thm. 2.2) *Assume (4.1), (4.3). Then $H = H_0 + \phi(v)$ is well defined as a form sum and yields a bounded below selfadjoint operator with $\text{Dom}|H|^{\frac{1}{2}} = \text{Dom}|H_0|^{\frac{1}{2}}$.*

The operator H defined as above is called an abstract Pauli-Fierz Hamiltonian.

4.2 Existence of a ground state for cutoff Hamiltonians

We introduce as in [BD] the infrared-cutoff objects

$$v_\sigma = F(\omega \geq \sigma)v, \quad H_\sigma = K \otimes \mathbb{1} + \mathbb{1} \otimes d\Gamma(\omega) + \phi(v_\sigma), \quad \sigma > 0,$$

where $F(\lambda \geq \sigma)$ denotes as usual a function of the form $\chi(\sigma^{-1}\lambda)$, where $\chi \in C^\infty(\mathbb{R})$, $\chi(\lambda) \equiv 0$ for $\lambda \leq 1$, $\chi(\lambda) \equiv 1$ for $\lambda \geq 2$.

An important step to prove that H has a ground state is to prove that H_σ has a ground state. The usual trick is to consider

$$\tilde{H}_\sigma = K \otimes \mathbb{1} + \mathbb{1} \otimes d\Gamma(\omega_\sigma) + \phi(v_\sigma),$$

where:

$$\omega_\sigma := F(\omega \leq \sigma)\sigma + (1 - F(\omega \leq \sigma))\omega = \omega + (\sigma - \omega)F(\omega \leq \sigma).$$

Note that since $\omega_\sigma \geq \sigma > 0$, \tilde{H}_σ is a massive Pauli-Fierz Hamiltonian. Moreover it is well known (see eg [G], [BD]) H_σ has a ground state iff \tilde{H}_σ does. The fact that \tilde{H}_σ has a ground state follows from an estimate on its essential spectrum (HVZ theorem). In [BD] this is shown using the condition that $\mathfrak{h} = L^2(\mathbb{R}^d, dk)$ and $\omega = \omega(k)$. Here we will replace this condition by the following more abstract condition, formulated using an additional selfadjoint operator $\langle x \rangle$ on \mathfrak{h} . Similar abstract conditions were introduced in [GP].

We will assume that there exists a selfadjoint operator $\langle x \rangle \geq 1$ on \mathfrak{h} such that the following conditions hold for all $\sigma > 0$:

$$(4.4) \quad \begin{aligned} & (i) (z - \langle x \rangle)^{-1} : \text{Dom} \omega_\sigma \rightarrow \text{Dom} \omega_\sigma, \forall z \in \mathbb{C} \setminus \mathbb{R}, \\ & (ii) [\langle x \rangle, \omega_\sigma] \text{ defined as a quadratic form on } \text{Dom} \langle x \rangle \cap \text{Dom} \omega \text{ is bounded,} \\ & (iii) \langle x \rangle^{-\epsilon} (\omega_\sigma + 1)^{-\epsilon} \text{ is compact on } \mathfrak{h} \text{ for some } 0 < \epsilon < \frac{1}{2}. \end{aligned}$$

The operator $\langle x \rangle$, called a *gauge*, is used to localize particles in \mathfrak{h} .

We assume also as in [BD]:

$$(4.5) \quad (K + 1)^{-\frac{1}{2}} \text{ is compact.}$$

This assumption means that the small system is confined.

Proposition 4.7 *Assume (4.1), (4.2), (4.3), (4.4), (4.5). Then*

$$\sigma_{\text{ess}}(\tilde{H}_\sigma) \subset [\inf \sigma(\tilde{H}_\sigma) + \sigma, +\infty[.$$

It follows that \tilde{H}_σ (and hence H_σ) has a ground state for all $\sigma > 0$.

Proof. By (4.3), $\phi(v_\sigma)$ is form bounded with respect to H_0 (and to $K \otimes \mathbb{1} + \mathbb{1} \otimes d\Gamma(\omega_\sigma)$) with the infinitesimal bound, hence $H_\sigma, \tilde{H}_\sigma$ are well defined as bounded below selfadjoint Hamiltonians.

We can follow the proof of [DG2, Thm. 4.1] or [GP, Thm. 7.1] for its abstract version. For ease of notation we denote simply \tilde{H}_σ by H , ω_σ by ω and v_σ by v . The key estimate is the fact that for $\chi \in C_0^\infty(\mathbb{R})$ one has

$$(4.6) \quad \chi(H^{\text{ext}})I^*(j^R) - I^*(j^R)\chi(H) \in o(1), \text{ when } R \rightarrow \infty.$$

(The extended operator H^{ext} and identification operator $I(j^R)$ are defined for example in [GP, Sect.2.4]). The two main ingredients of the proof of (4.6) are the estimates:

$$(4.7) \quad [F(\frac{\langle x \rangle}{R}), \omega_\sigma] \in O(R^{-1}), \quad F \in C_0^\infty(\mathbb{R}),$$

and

$$(4.8) \quad \omega_\sigma^{-\frac{1}{2}} F(\frac{\langle x \rangle}{R}) \geq 1) v_\sigma (K + 1)^{-\frac{1}{2}} \in o(R^0).$$

Now (4.8) follows from the fact that $v_\sigma (K + 1)^{-\frac{1}{2}}$ is compact (note that $\omega_\sigma^{-\frac{1}{2}}$ is bounded since $\omega_\sigma \geq \sigma$), and (4.7) follows from Lemma 4.8. The estimate (4.6) can then be proved exactly as in [GP, Lemma 6.3]. Note that here we prove only the \subset part of the HVZ theorem, which is sufficient for our purposes. The details are left to the reader. \square

Lemma 4.8 *Assume conditions (i), (ii) of (4.4). Then for all $F \in C_0^\infty(\mathbb{R})$ one has:*

$$\begin{aligned} F(\langle x \rangle) &: \text{Dom}\omega_\sigma \rightarrow \text{Dom}\omega_\sigma, \\ [F(\frac{\langle x \rangle}{R}), \omega_\sigma] &\in O(R^{-1}). \end{aligned}$$

Proof. The proof of the lemma is easy, using almost analytic extensions, as for example in [GP]. The details are left to the interested reader. \square

4.3 Existence of a ground state for massless models

Let us introduce the following hypothesis on the coupling operator ([BD, Hyp. F]):

$$(4.9) \quad \omega^{-1}v(K+1)^{-\frac{1}{2}} \text{ is compact.}$$

Theorem 4.9 *Assume (4.1), (4.2), (4.3), (4.4), (4.5) and (4.9). Then H has a ground state.*

Proof. we can follow the proof in [BD, Sect. 4]. The existence of ground state for H_σ ([BD, Prop. 4.5]) is shown in Prop. 4.7. The arguments in [BD, Sects 4.2, 4.3] based on the pullthrough and double pullthrough formulas are abstract and valid for any one particle operator ω . The only place where the fact that $\mathfrak{h} = L^2(\mathbb{R}^d, dk)$ and $\omega = \omega(k)$ appears is in [BD, Prop. 4.7] where the operator $|x| = |i\nabla_k|$ enters. In our situation it suffices to replace it by our gauge operator $\langle x \rangle$. The rest of the proof is unchanged. \square

4.4 Proof of Thm. 4.1

We now complete the proof of Thm. 4.1, by verifying the hypotheses of Thm. 4.9. We recall that $\mathfrak{h} = L^2(\mathbb{R}^d dx)$, $\omega = h^{\frac{1}{2}}$ and we will take $\langle x \rangle = (1 + x^2)^{\frac{1}{2}}$.

Proof of Thm. 4.1.

We saw in the proof of Lemma 3.3 that $v, \omega^{-\frac{1}{2}}v$ are bounded, hence in particular (4.2) is satisfied. By hypothesis (E2), $(K+1)^{-\frac{1}{2}}$ is compact, which implies that conditions (4.3) and (4.5) are satisfied.

We now check condition (4.4). Note that $\omega_\sigma = f(h)$ where $f \in C^\infty(\mathbb{R})$ with $f(\lambda) = \lambda^{\frac{1}{2}}$ for $\lambda \geq 2$. Clearly $\text{Dom}\omega_\sigma = H^1(\mathbb{R}^d)$ which is preserved by $(z - \langle x \rangle)^{-1}$, so (i) of (4.4) is satisfied. Condition (iii) is also obviously satisfied. It remains to check condition (ii). To this end we write $\omega_\sigma = f(h) = (h+1)g(h)$ where $g \in C^\infty(\mathbb{R})$ satisfies

$$g^{(n)}(\lambda) \in O(\langle \lambda \rangle^{-\frac{1}{2}-n}), \quad n \in \mathbb{N},$$

and hence

$$(4.10) \quad [\langle x \rangle, \omega_\sigma] = [\langle x \rangle, h]g(h) + (h+1)[\langle x \rangle, g(h)].$$

Since $\nabla a^{jk}(x), \nabla c(x), \nabla m(x)$ are bounded and $\text{Dom}h = H^2(\mathbb{R}^d)$ we see that

$$(4.11) \quad [\langle x \rangle, h](h+1)^{-\frac{1}{2}}, [[\langle x \rangle, h], h](h+1)^{-1} \text{ are bounded.}$$

In particular the first term in the r.h.s. of (4.10) is bounded. To estimate the second term, we use an almost analytic extension of g satisfying:

$$(4.12) \quad \begin{aligned} \tilde{g}|_{\mathbb{R}} = g, \quad & \left| \frac{\partial \tilde{g}}{\partial \bar{z}}(z) \right| \leq C_N \langle z \rangle^{-3/2-N} |\operatorname{Im} z|^N, \quad N \in \mathbb{N}, \\ & \operatorname{supp} \tilde{g} \subset \{z \in \mathbb{C} \mid |\operatorname{Im} z| \leq c(1 + |\operatorname{Re} z|)\}, \end{aligned}$$

(see eg [DG1, Prop. C.2.2]), and write

$$g(h) = \frac{i}{2\pi} \int_{\mathbb{C}} \frac{\partial \tilde{g}}{\partial \bar{z}}(z) (z - h)^{-1} dz \wedge d\bar{z}.$$

We perform a commutator expansion to obtain that:

$$[\langle x \rangle, g(h)] = g'(h)[\langle x \rangle, h] + R_2,$$

for

$$R_2 = \frac{i}{2\pi} \int_{\mathbb{C}} \frac{\partial \tilde{g}}{\partial \bar{z}}(z) (z - h)^{-2} [[\langle x \rangle, h]h](z - h)^{-1} dz \wedge d\bar{z}.$$

Since $|g'(\lambda)| \leq C \langle \lambda \rangle^{-3/2}$, $(h+1)g'(h)[\langle x \rangle, h]$ is bounded. To estimate the term $(h+1)R_2$, we use again (4.11) and the bound

$$\|(h+1)^\alpha (z-h)^{-1}\| \leq C \langle z \rangle^\alpha |\operatorname{Im} z|^{-1}, \quad \alpha = \frac{1}{2}, 1.$$

We obtain that

$$\|(h+1)R_2\| \leq C \| [[\langle x \rangle, h]h](h+1)^{-1} \| \int_{\mathbb{C}} \left| \frac{\partial \tilde{g}}{\partial \bar{z}}(z) \right| \langle z \rangle^2 |\operatorname{Im} z|^{-3} dz d\bar{z}.$$

This integral is convergent using the estimate (4.12). This completes the proof of (4.4).

It remains to check condition (4.9), i.e. the fact that the interaction is infrared regular. This is the only place where the lower bound on $m(x)$ enters. By Thm. A.8 we obtain that $\omega^{-3/2} \langle x \rangle^{-3/2-\epsilon}$ is bounded for all $\epsilon > 0$. By condition (E2), we obtain that $\langle X \rangle^{3/2+\epsilon} (K+1)^{-\frac{1}{2}}$ is bounded for all $\epsilon > 0$ small enough.

Therefore to check (4.9) it suffices to prove that the map

$$L^2(\mathbb{R}^3, dX) \ni u \mapsto \langle x \rangle^{3/2+\epsilon} \rho(x - X) \langle X \rangle^{-3/2-\epsilon} u(X) \in L^2(\mathbb{R}^3, dX) \otimes L^2(\mathbb{R}^3, dx)$$

is bounded, which is immediate since $\rho \in S(\mathbb{R}^3)$. This completes the proof of Thm. 4.1. \square

4.5 Existence of a ground state for non confined Hamiltonians

In this subsection we state the results on existence of a ground state if the electronic potential is not confining. As explained in the beginning of this section, one has to assume a *stability condition*, meaning that states near the bottom of the spectrum of H are confined in electronic variables from energy conservation arguments.

Definition 4.10 Let H be a Nelson Hamiltonian satisfying (E1), (B1). We assume for simplicity that the electronic potential $W(X)$ is bounded. Set for $R \geq 1$:

$$D_R = \{u \in \text{Dom}H \mid \mathbb{1}_{\{|X| \leq R\}}u = 0\}.$$

The ionization threshold of H is

$$\Sigma(H) := \lim_{R \rightarrow +\infty} \inf_{u \in D_R, \|u\|=1} (u|Hu).$$

The following theorem can easily be obtained by adapting the arguments in this section.

Theorem 4.11 Assume hypotheses (E1), (B1), $W \in L^\infty(\mathbb{R}^3)$ and $m(x) \geq a\langle x \rangle^{-1}$ for some $a > 0$. Then if the following stability condition is satisfied:

$$\Sigma(H) > \inf \sigma(H),$$

H has a ground state.

Sketch of proof. Assuming the stability condition one can prove using Agmon-type estimates as in [Gr] (see [P] for the case of the Nelson model) that if $\chi \in C_0^\infty(]-\infty, \Sigma(H)[$ then $e^{\beta|X|}\chi(H_\sigma)$ is bounded uniformly in $0 < \sigma \leq \sigma_0$ for σ_0 small enough. From this fact one deduces by the usual argument that H_σ has a ground state ψ_σ and that

$$(4.13) \quad \sup_{\sigma > 0} \|\langle X \rangle^N \psi_\sigma\| < \infty.$$

One can then follow the proof in [P, Thm. 1.2]. The key infrared regularity property replacing (4.9) is now

$$\sup_{\sigma > 0} \|\omega^{-1}v\psi_\sigma\|_{\mathcal{H} \otimes \mathfrak{h}} < \infty.$$

This estimate follows as in the proof of (4.9) from Thm. A.8 and the bound (4.13). The details are left to the reader. \square

5 Removal of the UV cutoff

Let us denote by $H(\rho)$ the Nelson Hamiltonian defined in (3.2) to emphasize its dependence on the charge density ρ . In [Ne] Nelson considered the limit of $H(\rho)$ for $\omega = (-\Delta + m^2)^{\frac{1}{2}}$ when ρ tends to the Dirac mass δ , leading to a interacting Hamiltonian with a *local* interaction. In this section we study the same problem for the Nelson model with variable coefficients.

In [Ne], the one-particle operator ω is diagonalized using the Fourier transform. In our case we will use instead the pseudodifferential calculus. We denote by $S^0(\mathbb{R}^3)$ the space:

$$S^0(\mathbb{R}^3) = \{f \in C^\infty(\mathbb{R}^3) \mid |\partial_x^\alpha f(x)| \leq C_\alpha, \alpha \in \mathbb{N}^3\}.$$

We will assume in addition to hypotheses (E1), (B1) that:

$$(N) \ A_{jk}(X), \ a_{jk}(x), \ c(x), \ m^2(x) \in S^0(\mathbb{R}^3).$$

It is easy to see that h can be rewritten as:

$$h = \sum_{jk} D_j c^{-2}(x) a^{jk}(x) D_j + v(x),$$

where $v \in S^0(\mathbb{R}^3)$, and that $c^{-2}(x) a^{jk}(x) \in S^0(\mathbb{R}^3)$. Changing notation, we will henceforth assume that

$$h = \sum_{jk} D_j a^{jk}(x) D_j + v(x),$$

where $[a_{jk}](x)$ satisfies (B1) and a^{jk} , $v \in S^0(\mathbb{R}^3)$.

5.1 Preparations

We refer the reader to Appendix B for the notation and for some background on pseudodifferential calculus. It will be useful later to consider $\omega = h^{\frac{1}{2}}$ as a pseudodifferential operator. Note first that

$$h = h^w(x, D_x),$$

for

$$h(x, \xi) = \sum_{1 \leq j, k \leq 3} \xi_j a^{jk}(x) \xi_k + c(x).$$

The symbol $h(x, \xi)$ belongs to $S(\langle \xi \rangle^2, g)$, for the standard metric $g = dx^2 + \langle \xi \rangle^{-2} d\xi^2$, and is elliptic in this class. By Lemma B.1 and Thm. B.3, we know that if $f \in S^p(\mathbb{R})$, then the operator $f(h)$ belongs to $\Psi^w(\langle \xi \rangle^{2p}, g)$.

If the model is massive, then picking a function $f \in S^{\frac{1}{2}}(\mathbb{R})$ equal to $\lambda^{\frac{1}{2}}$ in $\{\lambda \geq m/2\}$, we see that $\omega = f(h) \in \Psi^w(\langle \xi \rangle, g)$.

If the model is massless, we fix $\sigma > 0$ ($\sigma = 1$ will do) and pick $f \in C^\infty(\mathbb{R})$ such that:

$$f(\lambda) = \begin{cases} \lambda^{\frac{1}{2}} & \text{if } |\lambda| \geq 4\sigma^2, \\ \sigma & \text{if } |\lambda| \leq \sigma^2. \end{cases}$$

We set:

$$\omega_\sigma := f(h).$$

Again by Thm. B.3 we know that ω_σ belongs to $\Psi^w(\langle \xi \rangle, g)$. For simplicity in the massive case we set $\omega_\sigma := \omega$.

Consider now the operator:

$$T := K_0 \otimes \mathbb{1} + \mathbb{1} \otimes \omega_\sigma,$$

acting on $L^2(\mathbb{R}^3, dX) \otimes L^2(\mathbb{R}^3, dx)$. Clearly T is selfadjoint on its natural domain and $T \geq \sigma$.

Lemma 5.1 *Set*

$$M(\Xi, \xi) := \langle \Xi \rangle^2 + \langle \xi \rangle, \quad G = dX^2 + dx^2 + \langle \Xi \rangle^{-2} d\Xi^2 + \langle \xi \rangle^{-2} d\xi^2.$$

Then T^{-1} belongs to $\Psi^w(M^{-1}, G)$.

Proof. By Lemma B.2, the metric G and weight M satisfy all the conditions in Subsect. B.1. Clearly $T \in \Psi^w(M, G)$. We pick a function $f \in S^{-1}(\mathbb{R})$ such that $f(\lambda) = \lambda^{-1}$ in $\{\lambda \geq \sigma/2\}$. By Thm. B.3 $T^{-1} = f(T) \in \Psi^w(M^{-1}, G)$. \square

Let us fix another cutoff function $F(\lambda \geq \sigma) \in C^\infty(\mathbb{R})$ with

$$F(\lambda \geq \sigma) = \begin{cases} 1 & \text{for } |\lambda| \geq 4\sigma, \\ 0 & \text{for } |\lambda| \leq 2\sigma, \end{cases}$$

and set:

$$F(\lambda \leq \sigma) := 1 - F(\lambda \geq \sigma).$$

Lemma 5.2 *Set*

$$\beta(X, x) = \beta_X(x) := -T^{-1}F(\omega \geq \sigma)\omega^{-\frac{1}{2}}\rho_X = -T^{-1}F(\omega \geq \sigma)\omega_\sigma^{-\frac{1}{2}}\rho_X.$$

Then

(1) $\beta \in C^\infty(\mathbb{R}^6)$.

(2) Let $0 \leq \alpha < 1$. Then $\omega^\alpha \beta_X \in L^2(\mathbb{R}^3, dx)$ and there exists $s > 3/2$ such that

$$\|\omega^\alpha \beta_X\|_{L^2(\mathbb{R}^3, dx)} \leq C \|\rho\|_{H^{-s}(\mathbb{R}^3)},$$

uniformly in X .

(3) Let $\alpha > 0$. Then $\omega^{-\alpha} \nabla_X \beta_X \in L^2(\mathbb{R}^3, dx)$ and there exists $s > 3/2$ such that

$$\|\omega^{-\alpha} \nabla_X \beta_X\|_{L^2(\mathbb{R}^3, dx)} \leq C \|\rho\|_{H^{-s}(\mathbb{R}^3)},$$

uniformly in X .

(4) one has:

$$\omega^{-\frac{1}{2}}\rho_X + (K_0 \otimes \mathbb{1} + \mathbb{1} \otimes \omega)\beta_X = \omega^{-\frac{1}{2}}F(\omega \leq \sigma)\rho_X.$$

Proof. The function $\rho_X(x)$ is clearly C^∞ in (X, x) , so (1) follows from the fact that T^{-1} and $\omega_\sigma^{-\frac{1}{2}}F(\omega \geq \sigma)$ are pseudodifferential operators.

We claim that there exists a symbol $b_X(x, \xi) = b(X, x, \xi)$ such that

$$(5.1) \quad \begin{aligned} b(X, x, \xi) &\in S(\langle \xi \rangle^{-5/2}, dX^2 + dx^2 + \langle \xi \rangle^{-2} d\xi^2), \\ \beta_X &= b_X^{(1,0)}(x, D_x)\rho_X. \end{aligned}$$

Let us prove our claim. Applying Lemma 5.1 and (B.10), we know that $T^{-1} \in \Psi^{(1,0)}(M^{-1}, G)$. Setting $w(X, x) = T^{-1}\rho_X$, this yields:

$$\begin{aligned}
(5.2) \quad w(X, x) &= (2\pi)^{-3} \int e^{i(X \cdot \Xi + x \cdot \xi)} B(X, x, \Xi, \xi) \delta(\xi + \Xi) \hat{\rho}(\xi) d\xi d\Xi \\
&= (2\pi)^{-3} \int e^{i(x-X) \cdot \xi} B(X, x, -\xi, \xi) \hat{\rho}(\xi) d\xi \\
&= b_X^{(1,0)}(x, D_x) \rho_X
\end{aligned}$$

for

$$(5.3) \quad b_X(x, \xi) = B(X, x, -\xi, \xi),$$

where $B(X, x, \Xi, \xi) \in S(M^{-1}, G)$ is the $(1, 0)$ symbol of T^{-1} . This implies that:

$$b_X \in S(\langle \xi \rangle^{-2}, dX^2 + dx^2 + \langle \xi \rangle^{-2} d\xi^2).$$

Applying once again Thm. B.3, we know that $F(\omega \geq \sigma) \omega_\sigma^{-\frac{1}{2}} \in \Psi^{(1,0)}(\langle \xi \rangle^{-\frac{1}{2}}, g)$. By the composition property (B.11), we obtain our claim.

Statement (2) follows then from (5.1), if we note that $\omega^\alpha F(\omega \geq \sigma) \omega_\sigma^{-\frac{1}{2}} \in \Psi^{(1,0)}(\langle \xi \rangle^{\alpha-\frac{1}{2}}, g)$ and use the mapping property of pseudodifferential operators between Sobolev spaces recalled in (B.13). Statement (3) is proved similarly, using that

$$\nabla_X b_X(x, D_x) \rho_X = \partial_X b_X(x, D_x) \rho_X - b_X(x, D_x) \nabla_x \rho_X.$$

Finally (4) follows from the fact that $(\omega - \omega_\sigma) F(\omega \geq \sigma) = 0$. \square

5.2 Dressing transformation

Let ρ be a charge density as above. We set for $\kappa \gg 1$:

$$\rho^\kappa(x) := \kappa^3 \rho(\kappa x), \quad \rho_X^\kappa(x) = \rho^\kappa(x - X),$$

so that

$$(5.4) \quad \lim_{\kappa \rightarrow \infty} \rho_X^\kappa = q \delta_X \text{ in } H^{-s}(\mathbb{R}^3), \quad \forall s > 3/2.$$

This implies

$$(5.5) \quad \|\rho_X^\kappa\|_{H^{-s}(\mathbb{R}^3)} \leq C, \text{ uniformly in } X, \kappa, \text{ for all } s > 3/2.$$

We set

$$H^\kappa = H(\rho^\kappa),$$

and as in [Ne]:

$$U^\kappa := e^{i\phi(i\beta_X^\kappa)},$$

which is a unitary operator on \mathcal{H} . (Recall that β_X^κ is defined in Lemma 5.2).

Proposition 5.3 *Set*

$$\begin{aligned}
a_j^\kappa(X) &= \frac{1}{\sqrt{2}}a(\nabla_{X_j}\beta_X^\kappa), \\
R^\kappa &= 2\sum_{j,k}\nabla_{X_j}A_{jk}(X)a_k^\kappa(X) - a_j^{\kappa^*}(X)A_{jk}(X)\nabla_{X_k} \\
&\quad + \sum_{j,k}2a_j^{\kappa^*}(X)A_{jk}(X)a_k^\kappa(X) - a_j^{\kappa^*}(X)A_{jk}(X)a_k^{\kappa^*}(X) - a_j^\kappa(X)A_{jk}(X)a_k^\kappa(X), \\
V^\kappa(X) &= -(\rho_X^\kappa|\omega^{-1}F(\omega \geq \sigma)T^{-1}\rho_X^\kappa) + \frac{1}{2}(T^{-1}\rho_X^\kappa|F^2(\omega \geq \sigma)T^{-1}\rho_X^\kappa) \\
&\quad + \frac{1}{2}\sum_{jk}A_{jk}(X)(\nabla_{X_j}T^{-1}\rho_X^\kappa|\omega^{-1}F^2(\omega \geq \sigma)\nabla_{X_k}T^{-1}\rho_X^\kappa).
\end{aligned}$$

Then

$$\begin{aligned}
U^\kappa H^\kappa U^{\kappa^*} &= K + d\Gamma(\omega) + \phi(\omega^{-\frac{1}{2}}F(\omega \leq \sigma)\rho_X^\kappa) \\
&\quad + R^\kappa + V^\kappa(X).
\end{aligned}$$

Proof. We recall some well-known identities:

$$(5.6) \quad U^\kappa(d\Gamma(\omega) + \phi(\omega^{-\frac{1}{2}}\rho_{\kappa,X}))U^{\kappa^*} = d\Gamma(\omega) + \phi(\omega\beta_X^\kappa + \omega^{-\frac{1}{2}}\rho_X^\kappa) + \text{Re}(\frac{\omega}{2}\beta_X^\kappa + \omega^{-\frac{1}{2}}\rho_X^\kappa|\beta_X^\kappa).$$

Note that the scalar product in the rhs is real valued, since ρ_X^κ , β_X^κ and ω are real vectors and operators. Using once more that β_X^κ is real, we see that the operators $\phi(i\beta_X^\kappa)$ for different X commute, which yields:

$$U_\kappa D_{X_j} U^{\kappa^*} = D_{X_j} - \phi(i\nabla_{X_j}\beta_X^\kappa),$$

and hence:

$$U^\kappa K U^{\kappa^*} = \sum_{j,k} (D_{X_j} - \phi(i\nabla_{X_j}\beta_X^\kappa)) A_{jk}(X) (D_{X_k} - \phi(i\nabla_{X_k}\beta_X^\kappa)) + W(X).$$

We expand the squares in the r.h.s. using the definition of $a_j^\kappa(X)$ in the proposition. After rearranging the various terms, we obtain:

$$\begin{aligned}
U^\kappa K U^{\kappa^*} &= K + \phi(K_0\beta_X^\kappa) \\
&\quad + 2\sum_{j,k}\nabla_{X_j}A_{jk}(X)a_k^\kappa(X) - a_j^{\kappa^*}(X)A_{jk}(X)\nabla_{X_k} \\
&\quad + \sum_{j,k}2a_j^{\kappa^*}(X)A_{jk}(X)a_k^\kappa(X) - a_j^{\kappa^*}(X)A_{jk}(X)a_k^{\kappa^*}(X) - a_j^\kappa(X)A_{jk}(X)a_k^\kappa(X) \\
&\quad + \frac{1}{2}\sum_{jk}A_{jk}(X)(\nabla_{X_j}\beta_X^\kappa|\nabla_{X_k}\beta_X^\kappa).
\end{aligned}$$

This yields:

$$\begin{aligned}
U^\kappa H^\kappa U^{\kappa^*} &= K + d\Gamma(\omega) \\
&\quad + 2\sum_{j,k}\nabla_{X_j}A_{jk}(X)a_k^\kappa(X) - a_j^{\kappa^*}(X)A_{jk}(X)\nabla_{X_k} \\
&\quad + \sum_{j,k}2a_j^{\kappa^*}(X)A_{jk}(X)a_k^\kappa(X) - a_j^{\kappa^*}(X)A_{jk}(X)a_k^{\kappa^*}(X) - a_j^\kappa(X)A_{jk}(X)a_k^\kappa(X) \\
&\quad + \phi(\omega^{-\frac{1}{2}}\rho_X^\kappa + (K_0 + \omega)\beta_X^\kappa) \\
&\quad + (\omega^{-\frac{1}{2}}\rho_X^\kappa + \frac{1}{2}\omega\beta_X^\kappa|\beta_X^\kappa) + \frac{1}{2}\sum_{jk}A_{jk}(X)(\nabla_{X_j}\beta_X^\kappa|\nabla_{X_k}\beta_X^\kappa).
\end{aligned}$$

The sum of the second and third lines equals R^κ . By Lemma 5.2, the fourth line equals $\phi(\omega^{-\frac{1}{2}}F(\omega \leq \sigma)\rho_X)$. The fifth line equals $V^\kappa(X)$, using the definition of β_X . \square

5.3 Removal of the ultraviolet cutoff

Set

$$h_0(x, \xi) = \sum_{1 \leq j, k \leq 3} \xi_j a_{jk}(x) \xi_k, \quad K(X, \xi) = \sum_{1 \leq j, k \leq 3} \xi_j A_{jk}(X) \xi_k.$$

and:

$$(5.7) \quad E^\kappa(X) := -\frac{1}{2}(2\pi)^{-3} \int (h_0(X, \xi) + 1)^{-\frac{1}{2}} K(X, \xi) (K(X, \xi) + 1)^{-2} |\hat{\rho}|^2 (\xi \kappa^{-1}) d\xi.$$

Lemma 5.4 *Then there exists a bounded continuous potential V_{ren} such that:*

$$\lim_{\kappa \rightarrow +\infty} V^\kappa(X) - E^\kappa(X) = V_{\text{ren}}(X),$$

in $L^\infty(\mathbb{R}^3)$.

Theorem 5.5 *Assume hypotheses (E1), (B1), (N). Then the family of selfadjoint operators*

$$H^\kappa - E^\kappa(X)$$

converges in strong resolvent sense to a bounded below selfadjoint operator H^∞ .

Proof. By Prop. 5.6 below, $U^\kappa(H^\kappa - E^\kappa(X))U^{\kappa*}$ converges in norm resolvent sense to \hat{H}^∞ . Moreover by Lemma 5.2 (2), β_X^κ converges in $B(\mathcal{K}, \mathcal{K} \otimes \mathfrak{h})$ when $\kappa \rightarrow \infty$, hence U^κ converges strongly to some unitary operator U^∞ . It follows that H^κ converges in strong resolvent sense to

$$H^\infty = U^{\infty*} \hat{H}^\infty U^\infty. \quad \square$$

Proof of Lemma 5.4. For simplicity we will assume that the model is massive ($m > 0$), which allows to remove the cutoffs $F(\omega \geq \sigma)$ in the various formulas. The massless case can be treated similarly. Recall that:

$$(5.8) \quad \begin{aligned} T^{-1} \rho_X^\kappa &= b_X(x, D_x) \rho_X^\kappa, \\ \partial_X T^{-1} \rho_X^\kappa &= \partial_X b_X(x, D_x) \rho_X^\kappa - b_X(x, D_x) \partial_x \rho_X^\kappa, \end{aligned}$$

where $b_X(x, \xi)$ is defined in (5.3). Plugging the second identity in (5.8) into the formula giving $V_\kappa(X)$ we get:

$$V^\kappa(X) = V_1^\kappa(X) + V_2^\kappa(X),$$

for

$$\begin{aligned} V_1^\kappa(X) &= \frac{1}{2} \|b_X(x, D_x) \rho_X^\kappa\|^2 + \frac{1}{2} \sum_{jk} A_{jk}(X) (\partial_{X_j} b_X(x, D_x) \rho_X^\kappa | \omega^{-1} \partial_{X_k} b_X(x, D_x) \rho_X^\kappa) \\ &\quad - \sum_{jk} A_{jk}(X) (\partial_{X_j} b_X(x, D_x) \rho_X^\kappa | \omega^{-1} b_X(x, D_x) \partial_{x_k} \rho_X^\kappa), \\ V_2^\kappa(X) &= -(\rho_X^\kappa | \omega^{-1} b_X(x, D_x) \rho_X^\kappa) + \frac{1}{2} \sum_{jk} A_{jk}(X) (b_X(x, D_x) \partial_{x_j} \rho_X^\kappa | \omega^{-1} b_X(x, D_x) \partial_{x_k} \rho_X^\kappa). \end{aligned}$$

We will use that:

$$(5.9) \quad \begin{aligned} \rho_X^\kappa &\rightarrow q\delta_X \text{ in } H^s(\mathbb{R}^3), \quad \forall s < -\frac{3}{2}, \\ \partial_x \rho_X^\kappa &\rightarrow q\partial_x \delta_X \text{ in } H^s(\mathbb{R}^3), \quad \forall s < -\frac{5}{2}, \text{ uniformly in } X \in \mathbb{R}^3, \end{aligned}$$

where we recall that $q = \int_{\mathbb{R}^3} \rho(y) dy$. Using that $b_X(x, \xi) \in S(\langle \xi \rangle^{-2}, g)$ and the mapping properties of pseudodifferential operators between Sobolev spaces, we obtain that

$$\lim_{\kappa \rightarrow \infty} V_1^\kappa(X) =: V_1^\infty(X) \text{ exists uniformly for } X \in \mathbb{R}^3,$$

and $V_1^\infty(X)$ is a bounded continuous function, whose exact expression is obtained by replacing ρ_X^κ by $q\delta_X$ in the formula giving $V_1^\kappa(X)$.

We now consider the potential $V_2^\kappa(X)$, which will be seen to be logarithmically divergent when $\kappa \rightarrow \infty$. To extract its divergent part, we use symbolic calculus. We will use only the $(1, 0)$ quantization and omit the corresponding superscript. We first use Prop. B.4 for the metric G defined in Lemma 5.1. Note that the ‘Planck constant’ for the metric G is

$$\lambda(X, x, \Xi, \xi) = \min(\langle \Xi \rangle, \langle \xi \rangle).$$

Applying Prop. B.4, we obtain that the symbol $b_X(x, \xi)$ in (5.2) equals:

$$(5.10) \quad \begin{aligned} b_X(x, \xi) &= (K(X, \xi) + (h_0(x, \xi) + 1)^{\frac{1}{2}})^{-1} + S(\langle \xi \rangle^{-3}, g) \\ &= (K(X, \xi) + 1)^{-1} + S(\langle \xi \rangle^{-3}, g). \end{aligned}$$

The same argument for the metric g shows that $\omega^{-1} = d(x, D_x)$ for:

$$(5.11) \quad d(x, \xi) = (h_0(x, \xi) + 1)^{-\frac{1}{2}} + S(\langle \xi \rangle^{-2}, g).$$

Combining (5.10) and (5.11) we get that:

$$(5.12) \quad \begin{aligned} \omega^{-1} b_X(x, D_x) &= c_X(x, D_x) + r_X(x, D_x), \\ b_X^*(x, D_x) \omega^{-1} b_X(x, D_x) &= d_X(x, D_x) + s_X(x, D_x), \end{aligned}$$

where:

$$(5.13) \quad \begin{aligned} c_X(x, \xi) &= (h_0(x, \xi) + 1)^{-\frac{1}{2}} (K(X, \xi) + 1)^{-1}, \\ d_X(x, \xi) &= (h_0(x, \xi) + 1)^{-\frac{1}{2}} (K(X, \xi) + 1)^{-2}, \\ r_X(x, \xi) &\in S(\langle \xi \rangle^{-4}, g), \quad s_X(x, \xi) \in S(\langle \xi \rangle^{-6}, g), \text{ uniformly in } X \in \mathbb{R}^3. \end{aligned}$$

Setting

$$\tilde{V}_2^\kappa(X) = -(\rho_X^\kappa | c_X(x, D_x) \rho_X^\kappa) + \frac{1}{2} \sum_{jk} A_{jk}(X) (\partial_{x_j} \rho_X^\kappa | d_X(x, D_x) \partial_{x_k} \rho_X^\kappa),$$

we see using again (5.9) that

$$(5.14) \quad \lim_{\kappa \rightarrow \infty} V_2^\kappa(X) - \tilde{V}_2^\kappa(X) = V_2^\infty(X) \text{ exists uniformly for } X \in \mathbb{R}^3$$

and is a bounded continuous function. The potential $\tilde{V}_2^\kappa(X)$ can be explicitly evaluated. In fact:

$$(5.15) \quad \begin{aligned} & (\rho_X^\kappa | c_X(x, D_x) \rho_X^\kappa) \\ &= (2\pi)^{-3} \int e^{i(x-X) \cdot \xi} c_X(x, \xi) \rho_X^\kappa(x) \hat{\rho}(\kappa^{-1}\xi) dx d\xi \\ &= (2\pi)^{-3} \int e^{i(x-X) \cdot \xi} c_X(X, \xi) \rho_X^\kappa(x) \hat{\rho}(\kappa^{-1}\xi) dx d\xi + O(\kappa^{-1}) \log(\kappa) \\ &= (2\pi)^{-3} \int c_X(X, \xi) |\hat{\rho}|^2(\kappa^{-1}\xi) d\xi + O(\kappa^{-1}) \log(\kappa). \end{aligned}$$

Similarly

$$(5.16) \quad \begin{aligned} & (\partial_{x_j} \rho_X^\kappa | d_X(x, D_x) \partial_{x_k} \rho_X^\kappa) \\ &= (2\pi)^{-3} \int e^{i(x-X) \cdot \xi} \partial_j \rho_X^\kappa(x) d_X(x, \xi) i \xi_k \hat{\rho}(\kappa^{-1}\xi) dx d\xi \\ &= (2\pi)^{-3} \int e^{i(x-X) \cdot \xi} \rho_X^\kappa(x) d_X(x, \xi) \xi_j \xi_k \hat{\rho}(\kappa^{-1}\xi) dx d\xi \\ &\quad - (2\pi)^{-3} \int e^{i(x-X) \cdot \xi} \rho_X^\kappa(x) \partial_j d_X(x, \xi) i \xi_k \hat{\rho}(\kappa^{-1}\xi) dx d\xi. \end{aligned}$$

The second term in the rhs has a finite limit when $\kappa \rightarrow \infty$. By the same argument as above, we have:

$$(5.17) \quad \begin{aligned} & (2\pi)^{-3} \int e^{i(x-X) \cdot \xi} \rho_X^\kappa(x) d_X(x, \xi) \xi_j \xi_k \hat{\rho}(\kappa^{-1}\xi) dx d\xi \\ &= (2\pi)^{-3} \int e^{i(x-X) \cdot \xi} \rho_X^\kappa(x) d_X(X, \xi) \xi_j \xi_k \hat{\rho}(\kappa^{-1}\xi) dx d\xi + O(\kappa^{-1} \log(\kappa)) \\ &= (2\pi)^{-3} \int d_X(X, \xi) \xi_j \xi_k |\hat{\rho}|^2(\kappa^{-1}\xi) d\xi + O(\kappa^{-1} \log(\kappa)). \end{aligned}$$

Using the definition of $c_X(x, \xi)$ and $d_X(x, \xi)$ in (5.13), we get that

$$\begin{aligned} & -c_X(X, \xi) + \frac{1}{2} \sum_{jk} A_{jk}(X) \xi_j \xi_k d_X(X, \xi) \\ &= -\frac{1}{2} (h_0(X, \xi) + 1)^{-\frac{1}{2}} K(X, \xi) (K(X, \xi) + 1)^{-2}. \end{aligned}$$

Using the definition of $E^\kappa(X)$ and (5.15), (5.16) and (5.17) it follows that

$$\lim_{\kappa \rightarrow \infty} \tilde{V}_2^\kappa(X) - E^\kappa(X) \text{ exists uniformly for } X \in \mathbb{R}^3.$$

This completes the proof of the lemma. \square

Proposition 5.6 *Let*

$$\hat{H}^\kappa = U^\kappa H^\kappa U^{\kappa*} - E^\kappa(X).$$

Then there exists a bounded below selfadjoint operator \hat{H}^∞ such that

- (1) \hat{H}^κ converges to \hat{H}^∞ in norm resolvent sense;

$$(2) \quad D(|\hat{H}^\infty|^{\frac{1}{2}}) = D(H_0^{\frac{1}{2}}).$$

Proof. The proof is analogous to the one in [Ne], using Thm. B.6 so we will only sketch it. The important point is the convergence of R^κ as quadratic form on $D(|H_0|^{\frac{1}{2}})$ when $\kappa \rightarrow \infty$. The various terms in R^κ are estimated with the help of Lemma B.5, applied to the coupling operator: $v^\kappa = \nabla_{X_j} \beta_{X_j}^\kappa$. From Lemma 5.2 (3), we obtain that $\omega^{-\alpha} \nabla_{X_j} \beta_{X_j}^\kappa$ converges in $B(\mathcal{K}, \mathcal{K} \otimes \mathfrak{h})$ when $\kappa \rightarrow \infty$. The only remaining point to consider is the fact that powers of the number operator N appear in Lemma B.5. This is sufficient in the massive case since H_0 dominates N . In the massless case, we use the fact that $\beta_X^\kappa = F(\omega \geq \sigma/2) \beta_X^\kappa$. Therefore if we apply Lemma B.5, we can replace N by $d\Gamma(\mathbb{1}_{[\sigma/2, +\infty]}(\omega))$, which is dominated by H_0 . The rest of the proof is standard. \square

A Lower bounds for second order differential operators

In this section we prove various lower bounds for second order differential operators. These bounds are the key ingredient in the proof of the existence of a ground state for the Nelson model.

A.1 Second order differential operators

Let us introduce the class of second order differential operators that will be studied in this section. Let:

$$\begin{aligned} h_0 &= \sum_{1 \leq j, k \leq d} c(x)^{-1} D_j a^{jk}(x) D_k c(x)^{-1}, \\ h &= h_0 + v(x), \end{aligned}$$

with a^{jk} , c , v real functions and:

$$(A.1) \quad \begin{aligned} C_0 \mathbb{1} &\leq [a^{jk}(x)] \leq C_1 \mathbb{1}, \quad C_0 \leq c(x) \leq C_1, \quad C_0 > 0, \\ \partial_x^\alpha a^{jk}(x) &\in O(\langle x \rangle^{-1}), \quad |\alpha| \leq 1, \quad \partial_x^\alpha c(x) \in O(1), \quad |\alpha| \leq 2, \end{aligned}$$

$$(A.2) \quad v \in L^\infty(\mathbb{R}^d), \quad v \geq 0.$$

Clearly h_0 and h are selfadjoint and positive with domain $H^2(\mathbb{R}^d)$. We will always assume that $d \geq 3$.

A.2 Upper bounds on heat kernels

If K is a bounded operator on $L^2(\mathbb{R}^d, c^2 dx)$ we will denote by $K(x, y) \in \mathcal{D}'(\mathbb{R}^{2d})$ its distribution kernel. In this subsection we will prove the following theorem. We set:

$$\psi_\alpha(t, x) := \left(\frac{\langle x \rangle^2}{\langle x \rangle^2 + t} \right)^\alpha, \quad \alpha > 0.$$

Theorem A.1 *Assume in addition to (A.1), (A.2) that:*

$$v(x) \geq a\langle x \rangle^{-2}, \quad a > 0,$$

then there exists $C, c, \alpha > 0$ such that:

$$(A.3) \quad e^{-th}(x, y) \leq C\psi_\alpha(t, x)\psi_\alpha(t, y)e^{ct\Delta}(x, y), \quad \forall t > 0, \quad x, y \in \mathbb{R}^d.$$

If $c(x) \equiv 1$ or if h_0 is the Laplace-Beltrami operator for a Riemannian metric on \mathbb{R}^d , then Thm. A.1 is due to Zhang [Zh].

Remark A.2 *Conjugating by the unitary*

$$U : \begin{aligned} L^2(\mathbb{R}^d, dx) &\rightarrow L^2(\mathbb{R}^d, c^2(x)dx), \\ u &\mapsto c(x)^{-1}u, \end{aligned}$$

we obtain

$$\begin{aligned} \tilde{h}_0 &:= Uh_0U^{-1} = c(x)^{-2} \sum_{1 \leq j, k \leq d} D_j a^{jk}(x) D_k, \\ \tilde{h} &:= UhU^{-1} = \tilde{h}_0 + v(x), \end{aligned}$$

which are selfadjoint with domain $H^2(\mathbb{R}^d)$. Let $e^{-\tilde{h}}(x, y)$ for $t > 0$ the integral kernel of $e^{-\tilde{h}}$ i.e. such that

$$e^{-\tilde{h}}u(x) = \int_{\mathbb{R}^d} e^{-th}(x, y)u(y)c^2(y)dy, \quad t > 0.$$

Then since $e^{-th}(x, y) = c(x)e^{-\tilde{h}}(x, y)c(y)$, it suffices to prove Thm. A.1 for $e^{-\tilde{h}}$.

By the above remark, we will consider the operator \tilde{h}_0 (resp. \tilde{h}) and denote it again by h_0 (resp. h). We note that they are associated with the closed quadratic forms:

$$Q_0(f) = \int_{\mathbb{R}^d} \sum_{j, k} \partial_j \bar{f} a^{jk} \partial_k f \, dx, \quad Q(f) = Q_0(f) + \int_{\mathbb{R}^d} |f|^2 c^2 v \, dx,$$

with domain $H^1(\mathbb{R}^d)$.

Let us consider the semigroup $\{e^{-th}\}_{t \geq 0}$ generated by h . Since $\text{Dom}Q_0 = H^1(\mathbb{R}^d)$, we can apply [D, Thms. 1.3.2, 1.3.3] to obtain that e^{-th} is positivity preserving and extends as a semigroup of contractions on $L^p(\mathbb{R}^d, c^2 dx)$ for $1 \leq p \leq \infty$, strongly continuous on $L^p(\mathbb{R}^d, c^2 dx)$ if $p < \infty$. In other words $\{e^{-th}\}_{t \geq 0}$ is a Markov symmetric semigroup.

We first recall two results, taken from [PE] and [D].

Lemma A.3 *Assume (A.1), (A.2). Then there exist $c, C > 0$ such that:*

$$0 \leq e^{-th}(x, y) \leq Ce^{ct\Delta}(x, y), \quad \forall 0 < t, \quad x, y \in \mathbb{R}^d.$$

Proof. Since $v(x) \geq 0$ it follows from the Trotter-Kato formula that

$$0 \leq e^{-th}(x, y) \leq e^{-th_0}(x, y), \text{ a.e. } x, y.$$

The stated upper bound on $e^{-th_0}(x, y)$ is shown in [PE, Thm. 3.4]. \square

The following lemma is an extension of [D, Lemma 2.1.2] where the case $c(x) \equiv 1$ is considered.

Lemma A.4 *Assume (A.1), (A.2). Then:*

(1) e^{-th} is ultracontractive, i.e. e^{-th} is bounded from L^2 to L^∞ for all $t > 0$, and

$$c_t := \|e^{-th}\|_{L^2 \rightarrow L^\infty} = \sup_{f \in L^2} \frac{\|e^{-th}f\|_\infty}{\|f\|_2} \leq ct^{-d/4}$$

with some constant $c > 0$.

(2) e^{-th} is bounded from L^1 to L^∞ for all $t > 0$ and

$$\|e^{-th}\|_{L^1 \rightarrow L^\infty} \leq c_{t/2}^2.$$

(3) The kernel $e^{-th}(x, y)$ satisfies:

$$0 \leq e^{-th}(x, y) \leq c_{t/2}^2.$$

Proof. From Lemma A.3 we obtain that

$$\|e^{-th}f\|_\infty \leq C\|e^{ct\Delta}|f|\|_\infty \leq C't^{-d/4}\|f\|_2,$$

using the explicit form of the heat kernel of the Laplacian. This proves (1).

Taking adjoints we see that e^{-th} is also bounded from L^1 to L^2 with $\|e^{-th}\|_{L^1 \rightarrow L^2} \leq c_t$. It follows that

$$\|e^{-th}\|_{L^1 \rightarrow L^\infty} \leq \|e^{-th/2}\|_{L^2 \rightarrow L^\infty} \|e^{-th/2}\|_{L^1 \rightarrow L^2} \leq c_{t/2}^2,$$

which proves (2). Statement (3) is shown in [D, Lemma 2.1.2]. \square

We will deduce Thm. A.1 from the following result.

Theorem A.5 *Assume the hypotheses of Thm. A.1. Then there exists $C, \alpha > 0$ such that:*

$$e^{-th}(x, y) \leq Ct^{-d/2}\psi_\alpha(t, x)\psi_\alpha(t, y).$$

Proof of Theorem A.1:

Combining Lemma A.3 with Thm. A.5 we get:

$$\begin{aligned} e^{-th}(x, y) &= (e^{-th}(x, y))^\epsilon (e^{-th}(x, y))^{1-\epsilon} \\ &\leq Ct^{-\epsilon d/2} e^{-\epsilon(x-y)^2/2t} t^{-(1-\epsilon)d/2} \psi_\alpha(t, x)^{1-\epsilon} \psi_\alpha(t, y)^{1-\epsilon} \\ &\leq C't^{-d/2} e^{-c(x-y)^2/2t} \psi_\beta(t, x) \psi_\beta(t, y), \end{aligned}$$

for $\beta = (1 - \epsilon)\alpha$. This completes the proof of Thm. A.1. \square

It remains to prove Theorem A.5. To this end, we employ the following abstract result.

Lemma A.6 ([MS, Theorem B]) Let $(M, d\mu)$ be a locally compact measurable space with σ -finite measure μ and let A be a non-negative self-adjoint operator on $L^2(M, d\mu)$ such that

(i) $e^{-tA_1} := (e^{-tA}|_{L^1 \cap L^2})_{L^1 \rightarrow L^1}^{\text{clos}}$, $t \geq 0$ is a C_0 -semigroup of bounded operators, i.e.,

$$\|e^{-tA_1}\|_{L^1 \rightarrow L^1} \leq c_1, \quad t \geq 0.$$

(ii) e^{-tA} is bounded from L^1 to L^∞ with:

$$\|e^{-tA_1}\|_{L^1 \rightarrow L^\infty} \leq c_2 t^{-j}, \quad t > 0,$$

for some $j > 1$.

Assume moreover that there exists a family of weights $\psi(s, x)$ ($s > 0$) such that:

(B1) $\psi(s, x), \psi(s, x)^{-1} \in L^2_{\text{loc}}(M \setminus N, d\mu)$ for all $s > 0$, where N is a closed null set.

(B2) There is a constant \tilde{c} independent of s such that, for all $t \leq s$,

$$\|\psi(s, \cdot) e^{-tA} \psi(s, \cdot)^{-1} f\|_1 \leq \tilde{c} \|f\|_1, \quad f \in D_s,$$

where $D_s := \psi(s, \cdot) L^\infty_c(M \setminus N, d\mu)$

(B3) There exists $0 < \epsilon < 1$ and constants $\hat{c}_i > 0$, $i = 1, 2$ such that for any $s > 0$ there exists a measurable set $\Omega^s \subset M$ with

(a) $|\psi(s, x)|^{-\epsilon} \leq \hat{c}_1$ for all $x \in M \setminus \Omega^s$,

(b) $|\psi(s, x)|^{-\epsilon} \in L^q(\Omega^s)$ and $\|\psi(s, \cdot)^{-\epsilon}\|_{L^q(\Omega^s)} \leq \hat{c}_2 s^{j/q}$ with $q = 2/(1 - \epsilon)$ and $j > 1$ is the exponent in condition (ii).

Then there is a constant C such that

$$|e^{-tA}(x, y)| \leq C t^{-j} |\psi(t, x) \psi(t, y)|, \quad \forall t > 0, \text{ a.e. } x, y \in M.$$

To verify condition (B2) of Lemma A.6, we will use the following lemma.

Lemma A.7 ([MS, Criterion 2]) Let e^{-tA} be a C_0 -semigroup on $L^2(M, d\mu)$. Denote by $\langle \cdot, \cdot \rangle$ the scalar product on $L^2(M, d\mu)$. Then:

$$\|e^{-tA} f\|_{L^\infty} \leq \|f\|_{L^\infty}, \quad f \in L^2 \cap L^\infty, \quad t > 0,$$

if and only if:

$$(A.4) \quad \text{Re} \langle f - f_\wedge, Af \rangle \geq 0, \quad f \in D(A),$$

where $f_\wedge = (|f| \wedge 1) \text{sgn} f$ with $\text{sgn} f(x) := f(x)/|f(x)|$ if $|f(x)| \neq 0$ and $\text{sgn} f(x) = 0$ if $f(x) = 0$.

Proof of Thm. A.5: We will prove that there exists $\alpha > 0$ such that the hypotheses of Lemma A.6 are satisfied for $(M, d\mu) = (\mathbb{R}^d, c^2(x) dx)$, $A = h$ and $\psi(s, x) = \psi_\alpha(s, x)$. For ease of notation we will often denote ψ_α simply by ψ .

From the discussion before Lemma A.4, we know that e^{-th} extends as a C_0 -semigroup of contractions of $L^1(\mathbb{R}^d, c^2 dx)$, which implies that hypothesis (i) holds with $c_1 = 1$. Hypothesis (ii) with $j = d/2$ follows from (2) of Lemma A.4. Note that $d/2 > 1$ since $d \geq 3$.

We now check that conditions (B) are satisfied by ψ_α provided we choose $\alpha = \alpha_0 a^{\frac{1}{2}}$ for some constant α_0 . Since ψ, ψ^{-1} are bounded, condition (B1) is satisfied for all $\alpha > 0$. Set $\Omega^s := \{x \in \mathbb{R}^d \mid \langle x \rangle^2 \leq s\}$. Then

$$\psi(x)^{-\epsilon} = \left[\frac{\langle x \rangle^2 + s}{\langle x \rangle^2} \right]^{\alpha\epsilon} \leq 2^{\alpha\epsilon}, \quad \forall x \notin \Omega^s,$$

which proves the bound (a) of (B3) for all $\alpha > 0$. Take now $0 < \epsilon < \frac{d}{d+4\alpha}$ so that we see that $d - 2\alpha\epsilon q > 0$ for $q = 2/(1 - \epsilon)$. If $0 \leq s < 1$ $\Omega^s = \emptyset$ and (b) of (B3) is satisfied. If $s \geq 1$ we have:

$$\begin{aligned} \|\psi^{-\epsilon}\|_{L^q(\Omega^s)}^q &= \int_{\Omega^s} \left[\frac{\langle x \rangle^2 + s}{\langle x \rangle^2} \right]^{\alpha\epsilon q} c^2(x) dx \\ &\leq C_1^2 (2s)^{\alpha\epsilon q} \int_{\{|x| \leq \sqrt{s}\}} |x|^{-2\alpha\epsilon q} dx \\ &= C s^{\alpha\epsilon q} \int_0^{\sqrt{s}} r^{d-2\alpha\epsilon q-1} dr = C' s^{d/2}. \end{aligned}$$

Hence (b) is satisfied for $j = d/2$.

It remains to check (B2). To avoid confusion, we denote by $\langle g, f \rangle$ the scalar product in $L^2(\mathbb{R}^d, c^2(x) dx)$ and by $(g|f)$ the usual scalar product in $L^2(\mathbb{R}^d, dx)$.

Since ψ, ψ^{-1} are C^∞ and bounded with all derivatives, we see that $\{\psi e^{-th} \psi^{-1}\}_{t \geq 0}$ is a C_0 -semigroup on $L^2(\mathbb{R}^d, c^2 dx)$, with generator

$$h_\psi := \psi h \psi^{-1}, \quad \text{Dom } h_\psi = H^2(\mathbb{R}^d).$$

We claim that there exists $\alpha > 0$ such that

$$(A.5) \quad \|e^{-th_\psi}\|_{L^1 \rightarrow L^1} \leq C, \quad \text{uniformly for } 0 \leq t \leq s.$$

By duality, (A.5) will follow from (A.6):

$$(A.6) \quad \|e^{-th_\psi^*}\|_{L^\infty \rightarrow L^\infty} \leq C, \quad \text{uniformly for } 0 \leq t \leq s.$$

To prove (A.6), we will apply Lemma A.7. To avoid confusion, $\partial_j f(x)$ will denote a partial derivative of the function f , while $\nabla_j f(x)$ denote the product of the operator ∇_j and the operator of multiplication by the function f .

Setting $b_i = \psi^{-1}\partial_i\psi$, we have:

$$\begin{aligned}
h_\psi^* &= \psi^{-1}h\psi \\
&= -c(x)^{-2} \sum_{j,k} \nabla_j a^{jk}(x) \nabla_k - \sum_{j,k} c^{-2}(x) b_j(x) a^{jk}(x) \nabla_k - c^{-2}(x) \nabla_j a^{jk}(x) b_k(x) \\
&\quad + v(x) - c^{-2}(x) \sum_{j,k} b_j(x) a^{jk}(x) b_k(x) \\
&= -c(x)^{-2} \sum_{j,k} \nabla_j a^{jk}(x) \nabla_k - 2c(x)^{-2} \sum_{j,k} b_j(x) a^{jk}(x) \nabla_k + w(x),
\end{aligned}$$

where:

$$w(x) = v(x) - c(x)^{-2} \sum_{j,k} b_j(x) a^{jk}(x) b_k(x) - c(x)^{-2} \sum_{j,k} a^{jk}(x) \partial_j b_k(x) - c(x)^{-2} \sum_{j,k} (\partial_j a^{jk})(x) b_k(x).$$

Clearly $\text{Dom} h_\psi^* = H^2(\mathbb{R}^d)$. To simplify notation, we set $A(x) = [a^{jk}(x)]$, $F(x) = (b_1(x), \dots, b_d(x))$. The identity above becomes:

$$\begin{aligned}
(A.7) \quad h_\psi^* &= -c^{-2} \nabla_x A \nabla_x - c^{-2} F A \nabla_x - c^{-2} \nabla_x A F + v - c^{-2} F A F, \\
&= -c^{-2} \nabla_x A \nabla_x - 2c^{-2} F A \nabla_x + w.
\end{aligned}$$

We note that $b_j(x) = \alpha s x_j \langle x \rangle^{-2} (\langle x \rangle^2 + s)^{-1}$, which implies that:

$$|b_j(x)| \leq C \alpha \langle x \rangle^{-1}, \quad |\nabla_x b_j(x)| \leq C \alpha \langle x \rangle^{-2}, \text{ for some } C > 0.$$

Since $v(x) \geq a \langle x \rangle^{-2}$, this implies using also (A.1) that:

$$(A.8) \quad v(x) - c(x)^{-2} F A F(x) \geq 0, \quad w(x) \geq 0,$$

for $\alpha > 0$ small enough.

This implies that

$$(A.9) \quad \text{Re} \langle f, h_\psi^* f \rangle = -(\nabla_x f |A \nabla_x f) + (f | (c^2 v - F A F) f) \geq 0, \quad \text{for } f \in H^1(\mathbb{R}^d).$$

It follows that h_ψ^* is maximal accretive, hence $e^{-th_\psi^*}$ is a C_0 -semigroup of contractions by the Hille-Yosida theorem.

To check condition (A.4) in Lemma A.7 we follow [MS], with some easy modifications. We write

$$f - f_\Lambda = \text{sgn} f \chi, \quad \chi := \mathbb{1}_{\{|f| \geq 1\}} (|f| - 1),$$

and note that if $f \in \text{Dom} h_\psi^* \subset H^1(\mathbb{R}^d)$ then $|f|$, $\text{sgn} f$, $\chi \in H^1(\mathbb{R}^d)$ with

$$(A.10) \quad \nabla \text{sgn} f = \frac{\nabla f}{|f|} - f \frac{\nabla f}{|f|^2}, \quad \nabla \chi = \mathbb{1}_{\{|f| \geq 1\}} \nabla |f|, \quad \nabla |f| = \frac{1}{2|f|} (\bar{f} \nabla f + f \nabla \bar{f}).$$

We have:

$$\begin{aligned}
\langle f - f_\Lambda, h_\psi^* f \rangle &= (\nabla(f - f_\Lambda) |A \nabla f) - 2(F(f - f_\Lambda) |A \nabla f) + ((f - f_\Lambda) | c^2 w f) \\
&=: C_1(f) + C_2(f) + C_3(f).
\end{aligned}$$

Using (A.10), we have:

$$\begin{aligned}
C_1(f) &= (\nabla(f - f_\Lambda)|A\nabla f) \\
&= (\nabla f|\frac{\chi}{|f|}A\nabla f) - (\nabla|f||\bar{f}\frac{\chi}{|f|^2}A\nabla f) + (\nabla\chi|\frac{\bar{f}}{|f|}A\nabla f) \\
&=: B_1(f) + B_2(f) + B_3(f).
\end{aligned}$$

Clearly $B_1(f)$ is real valued. Next:

$$(A.11) \quad \operatorname{Re}B_2(f) = -\frac{1}{2}(\nabla|f||\frac{\chi}{|f|^2}A(\bar{f}\nabla f + f\nabla\bar{f})) = -(\nabla|f||\frac{\chi}{|f|}A\nabla|f|),$$

using (A.10). Similarly:

$$(A.12) \quad \operatorname{Re}B_3(f) = \frac{1}{2}(\nabla\chi|\frac{1}{|f|}A(\bar{f}\nabla f + f\nabla\bar{f})) = (\nabla\chi|A\nabla\chi),$$

using again (A.10). We estimate now $\operatorname{Re}C_2(f)$. We have:

$$(A.13) \quad \operatorname{Re}C_2(f) = -2\operatorname{Re}(F(f - f_\Lambda)|A\nabla f) = \frac{1}{2}(\chi|\frac{F}{|f|}A(\bar{f}\nabla f + f\nabla\bar{f})) = -2(F\chi|A\nabla\chi).$$

We estimate now $\operatorname{Re}C_3(f)$. We have:

$$(A.14) \quad \operatorname{Re}C_3(f) = \operatorname{Re}(f - f_\Lambda|c^2wf) = \operatorname{Re}(\chi|c^2w|f|) = (\chi|c^2w|f|) = (\chi|c^2w\chi) + (\chi|c^2w).$$

Collecting (A.11) to (A.13), we obtain that:

$$\begin{aligned}
(A.15) \quad \operatorname{Re}\langle f - f_\Lambda, h_\psi^* f \rangle &= (\nabla f|\frac{\chi}{|f|}A\nabla f) - (\nabla|f||\frac{\chi}{|f|}A\nabla|f|) \\
&\quad + (\nabla\chi|A\nabla\chi) - 2(F\chi|A\nabla\chi) + (\chi|c^2w\chi) \\
&\quad + (\chi|c^2w).
\end{aligned}$$

We use now the pointwise identity:

$$\begin{aligned}
&\nabla\bar{f}A\nabla f - \nabla|f|A\nabla|f| \\
&= \nabla\bar{f}A\nabla f - \frac{1}{4|f|^2}(\bar{f}\nabla f + f\nabla\bar{f})A(\bar{f}\nabla f + f\nabla\bar{f}) \\
&= \frac{1}{4|f|^2}(2|f|^2\nabla\bar{f}A\nabla f - f^2\nabla\bar{f}A\nabla\bar{f} - \bar{f}^2\nabla fA\nabla f) \\
&= \frac{1}{|f|^2}(\operatorname{Re}f\nabla\operatorname{Im}f - \operatorname{Im}f\nabla\operatorname{Re}f)A(\operatorname{Re}f\nabla\operatorname{Im}f - \operatorname{Im}f\nabla\operatorname{Re}f) \geq 0.
\end{aligned}$$

Hence the first line in the rhs of (A.15) is positive. Concerning the third line, we recall that (A.8) implies that $w \geq 0$ if $\alpha = \alpha_0 a$. Since $\chi \geq 0$ the third line is also positive. Therefore:

$$\begin{aligned}
\operatorname{Re}\langle f - f_\Lambda, h_\psi^* f \rangle &\geq (\nabla\chi|A\nabla\chi) - 2(F\chi|A\nabla\chi) + (\chi|c^2w\chi) \\
&= \langle \chi, h_\psi^* \chi \rangle = \operatorname{Re}\langle \chi, h_\psi^* \chi \rangle,
\end{aligned}$$

using (A.7) and the fact that χ is real. Using (A.9) we obtain condition (A.4). This completes the proof of Thm. A.5. \square

A.3 Lower bounds for differential operators

We now deduce lower bounds for powers of h from the heat kernel bounds in Subject. A.2.

Theorem A.8 *Assume hypotheses (A.1), (A.2) and*

$$v(x) \geq a\langle x \rangle^{-2}, \quad a > 0.$$

Then

$$h^{-\beta} \leq C\langle x \rangle^{2\beta+\epsilon}, \quad \forall 0 \leq \beta \leq d/2, \quad \epsilon > 0.$$

We start by an easy consequence of Sobolev inequality.

Lemma A.9 *On $L^2(\mathbb{R}^d)$ the following inequality holds:*

$$(-\Delta)^{-\gamma} \leq C\langle x \rangle^{2\delta}, \quad \forall 0 \leq \gamma < d/2, \quad \delta > \gamma.$$

Proof. We have

$$(f|(-\Delta)^{-\gamma}f) = C \int \int \frac{\bar{f}(x)f(y)}{|x-y|^{d-2\gamma}} dx dy, \quad \forall 0 < \gamma < n/2.$$

By the Sobolev inequality ([RS2, Equ. IX.19]):

$$\int \int \frac{\bar{f}(x)f(y)}{|x-y|^{d-2\gamma}} dx dy \leq C\|f\|_r^2,$$

for $r = 2d/(d + 2\gamma)$. We write then $f = \langle x \rangle^{-\alpha} \langle x \rangle^\alpha f$ and use Hölder inequality to get:

$$\|f\|_r \leq \|\langle x \rangle^{-\alpha}\|_p \|\langle x \rangle^\alpha f\|_q, \quad p^{-1} + q^{-1} = r^{-1}.$$

We choose $q = 2, p = d/\gamma$. The function $\langle x \rangle^{-\alpha}$ belongs to $L^{d/\gamma}$ if $\alpha > \gamma$. This implies the lemma. \square

Proof of Thm. A.8.

We first recall the formula:

$$(A.16) \quad \lambda^{-1-\nu} = \frac{1}{\Gamma(\nu+1)} \int_0^{+\infty} e^{-t\lambda} t^\nu dt, \quad \nu > -1.$$

In the estimates below, various quantities like $(f|h^{-\delta}f)$ appear. To avoid domain questions, it suffices to replace h by $h + m$, $m > 0$, obtaining estimates uniform in m and letting $m \rightarrow 0$ at the end of the proof. We will hence prove the bounds

$$(A.17) \quad (f|(h+m)^{-\beta}f) \leq C(f|\langle x \rangle^{2\beta+\epsilon}f), \quad \forall f \in C_0^\infty(\mathbb{R}^d),$$

uniformly in $m > 0$. Moreover we note that it suffices to prove (A.17) for $f \geq 0$. In fact it follows from (A.16) that $(h+m)^{-\beta}$ has a positive kernel. Therefore

$$(f|(h+m)^{-\beta}f) \leq (|f|(h+m)^\beta|f) \leq C(|f|\langle x \rangle^{2\beta+\epsilon}|f|) = C(f|\langle x \rangle^{2\beta+\epsilon}f),$$

and (A.17) extends to all $f \in C_0^\infty(\mathbb{R}^d)$.

We will use the bound (A.3) in Thm. A.1, noting that if (A.3) holds for some $\alpha_0 > 0$ it holds also for all $0 < \alpha \leq \alpha_0$. We use the inequality

$$\left(\frac{\langle x \rangle^2}{\langle x \rangle^2 + t} \right) \left(\frac{\langle y \rangle^2}{\langle y \rangle^2 + t} \right) \leq \frac{\langle y \rangle^2}{t},$$

and get for $f \in C_0^\infty(\mathbb{R}^d)$, $f \geq 0$:

$$\begin{aligned} h^{-\beta} f(x) &= c \int_0^{+\infty} t^{\beta-1} e^{-th} f(x) dt \\ &\leq C \int_0^{+\infty} t^{\beta-\alpha-1} (e^{ct\Delta} \langle x \rangle^{2\alpha}) f(x) dt \\ &= C' (-\Delta)^{\beta-\alpha} \langle x \rangle^{2\alpha} f(x), \end{aligned}$$

as long as $\beta > \alpha$, using again (A.16). Integrating this pointwise inequality, we get that

$$(f|h^{-2\beta} f) \leq C(f|\langle x \rangle^{2\alpha} (-\Delta)^{-2(\beta-\alpha)} \langle x \rangle^{2\alpha} f).$$

We can apply Lemma A.9 as long as $2(\beta - \alpha) < d/2$, and obtain

$$(f|h^{-2\beta} f) \leq C(f|\langle x \rangle^{4\beta+\epsilon} f), \quad \forall \epsilon > 0,$$

if $\alpha < \beta < \alpha + d/4$. Since α can be taken arbitrarily close to 0, this completes the proof of the theorem. \square

B Background on pseudodifferential calculus

In this section we recall various standard results on pseudodifferential calculus that will be needed in the sequel. It is convenient to use the language of the Weyl-Hörmander calculus.

B.1 Symbol classes

We start by recalling the definition of symbol classes and weights. Let g be a Riemannian metric on \mathbb{R}^d , i.e. a map:

$$g : \mathbb{R}^d \ni X \mapsto g_X,$$

with values in positive definite quadratic forms on \mathbb{R}^d . If $M : \mathbb{R}^d \rightarrow]0, +\infty[$ is a strictly positive function called a *weight*, one denotes by $S(M, g)$ the symbol class of functions in $C^\infty(\mathbb{R}^d)$ such that:

$$\left| \prod_{i=1}^k (v_i \cdot \nabla_X) a(X) \right| \leq C_k M(X) \prod_{i=1}^k |g_X(v_i)|^{\frac{1}{2}},$$

uniformly for $X \in \mathbb{R}^d$, $v_1, \dots, v_k \in \mathbb{R}^d$ and $k \in \mathbb{N}$. The best constants C_k are seminorms on $S(M, g)$.

Usually $d = 2n$ and one sets $\mathbb{R}^d \ni X = (x, \xi) \in \mathbb{R}^n \times \mathbb{R}^n$. If

$$(B.1) \quad g_X = dx^2 + \langle \xi \rangle^{-2} d\xi^2$$

and $M(X) = \langle \xi \rangle^m$, the symbol class $S(M, g)$ is the usual symbol class

$$S_{1,0}^m = \{a : |\partial_x^\alpha \partial_\xi^\beta a(x, \xi)| \leq C_{\alpha,\beta} \langle \xi \rangle^{m-|\beta|}, \alpha, \beta \in \mathbb{N}^n\}.$$

For simplicity we will also denote by $S^p(\mathbb{R})$, $p \in \mathbb{R}$, the space

$$(B.2) \quad S^p(\mathbb{R}) = \{f : |f^{(k)}(\lambda)| \leq C_k \langle \lambda \rangle^{p-k}, k \in \mathbb{N}\},$$

ie $S^p(\mathbb{R}) = S(\langle \lambda \rangle^p, \langle \lambda \rangle^{-2} d\lambda^2)$.

If one equips \mathbb{R}^{2n} with the usual symplectic form σ , one can consider the dual metric g_X^σ . Diagonalising g_X in (linear) symplectic coordinates on \mathbb{R}^{2n} , one can write:

$$g_X(dx, d\xi) = \sum_{i=1}^n \frac{dx_i^2}{a_i^2(X)} + \frac{d\xi_i^2}{\alpha_i^2(X)},$$

and

$$g_X^\sigma(dx, d\xi) = \sum_{i=1}^n \alpha_i^2(X) dx_i^2 + a_i^2(X) d\xi_i^2.$$

One introduces also the two functions $\lambda(X)$, $\Lambda(X)$ which are the best functions such that

$$\lambda(X)^2 g_X \leq g_X^\sigma \leq \Lambda(X)^2 g_X,$$

equal to:

$$\lambda(X) = \min_i a_i(X) \alpha_i(X), \quad \Lambda(X) = \max_i a_i(X) \alpha_i(X).$$

The function $\lambda(X)$ plays the role of the Planck constant.

One says that g is a *Hörmander metric*, if the following conditions are satisfied:

- (1) *uncertainty principle*: $\lambda(X) \geq 1$;
- (2) *slowness*: there exists $C > 0$ such that

$$(B.3) \quad g_Y(X - Y) \leq C^{-1} \Rightarrow (g_Y(\cdot)/g_X(\cdot))^{\pm 1} \leq C;$$

- (3) *temperateness*: there exist $C > 0$, $N \in \mathbb{N}$ such that:

$$(B.4) \quad (g_Y(\cdot)/g_X(\cdot))^{\pm 1} \leq C (1 + g_Y^\sigma(Y - X))^N.$$

One says that a weight M is *admissible* for g if there exist $C > 0$, $N \in \mathbb{N}$ such that:

$$(B.5) \quad (M(Y)/M(X))^{\pm 1} \leq \begin{cases} C, & \text{for } g_Y(X - Y) \leq C^{-1}, \\ C(1 + g_Y^\sigma(X - Y))^N, & \text{for } X, Y \in \mathbb{R}^{2n}. \end{cases}$$

The metric g is *geodesically temperate* if g is temperate and if there exist $C > 0$ and $N \in \mathbb{N}$ such that:

$$(B.6) \quad (g_Y(\cdot)/g_X(\cdot))^{\pm 1} \leq C(1 + d^\sigma(X, Y))^N,$$

where d^σ is the geodesic distance for the metric g^σ .

The metric g is *strongly slow* if there exists $C > 0$ such that:

$$(B.7) \quad g_Y^\sigma(X - Y) \leq C^{-1}\Lambda(Y)^2 \Rightarrow (g_Y(\cdot)/g_X(\cdot))^{\pm 1} \leq C.$$

Lemma B.1 *The metric $dx^2 + \langle \xi \rangle^{-2}d\xi^2$ and weight $\langle \xi \rangle^\alpha$ for $\alpha \in \mathbb{R}$ satisfy all the above conditions.*

Proof. Most conditions are immediate, except the last two. To check (B.6), we note that $d^\sigma(X, Y) \leq |\xi - \eta|$, from which (B.6) follows. (B.7) follows from the fact that $\Lambda(X) = \langle \xi \rangle$. \square

Lemma B.2 *Assume that (g_i, M_i) , $i = 1, 2$ are two metrics and weights on \mathbb{R}^{2n_i} satisfying all the above conditions. Then (g, M) on \mathbb{R}^{2n} satisfy all the above conditions for $n = n_1 + n_2$ and:*

$$g_X(dx) = g_{X_1}(dx_1) + g_{X_2}(dx_2), \quad M(X) = M_1(X_1) + M_2(X_2).$$

B.2 Pseudodifferential calculus

To a symbol $a \in S'(\mathbb{R}^{2n})$, one can associate the operator defined by:

$$(B.8) \quad a^w(x, D)u(x) = (2\pi)^{-n} \int e^{i(x-y)\cdot\xi} a\left(\frac{x+y}{2}, \xi\right) u(y) dy d\xi,$$

called the *Weyl quantization* of a , which is well defined as a bounded operator from $S(\mathbb{R}^n)$ into $S'(\mathbb{R}^n)$. Let (g, M) be a metric and weight satisfying (B.3), (B.4), (B.5). We set:

$$\Psi^w(M, g) = \{a^w : a \in S(M, g)\}.$$

If $a \in S(M, g)$ then $\text{Op}^w(a)$ sends $S(\mathbb{R}^n)$ into itself. Moreover as quadratic forms on $S(\mathbb{R}^n)$:

$$(a^w)^* = \bar{a}^w.$$

One often uses also the the $(1, 0)$ quantization defined by:

$$(B.9) \quad a^{1,0}(x, D)u(x) = (2\pi)^{-n} \int e^{i(x-y)\cdot\xi} a(x, \xi) u(y) dy d\xi.$$

One has with obvious notations:

$$(B.10) \quad \Psi^w(M, g) = \Psi^{(1,0)}(M, g),$$

Moreover

$$(B.11) \quad \Psi^\#(M_1, g) \times \Psi^\#(M_2, g) \subset \Psi^\#(M_1 M_2, g),$$

where $\# = w$ or $(1, 0)$ and if $a \in S(M, g)$

$$(B.12) \quad a^w(x, D_x) = b^{(1,0)}(x, D_x), \text{ where } a - b \in S(M\lambda^{-1}, g).$$

Let now g be the standard metric defined in (B.1) and $H^s(\mathbb{R}^d)$ be the Sobolev space of order $s \in \mathbb{R}$. Then

$$(B.13) \quad \Psi^\#(\langle \xi \rangle^p, g) \subset B(H^s(\mathbb{R}^d), H^{s-p}(\mathbb{R}^d)),$$

and the norm of $a^\#$ in $B(H^s(\mathbb{R}^d), H^{s-p}(\mathbb{R}^d))$ is controlled by a finite number of seminorms of a in $S(\langle \xi \rangle^p, g)$.

B.3 Functional calculus for pseudodifferential operators

Assume that the weight M satisfies:

$$(B.14) \quad M(X) \leq C(1 + \lambda(X))^N, \quad C > 0, \quad N \in \mathbb{N}.$$

A symbol $a \in S(M, g)$ is *elliptic* if

$$(B.15) \quad 1 + |a(X)| \geq C^{-1}M(X).$$

The following theorem is shown in [Bo, Corr. 4.5]:

Theorem B.3 *Assume that (M, g) satisfy all the conditions in Subsect. B.1. Assume moreover that $M \geq 1$, $a \in S(M, g)$ is real and elliptic, and a^w is essentially selfadjoint on $S(\mathbb{R}^n)$. Then if $f \in S^p(\mathbb{R})$, the operator $f(a^w)$ belongs to $\Psi^w(M^p, g)$.*

The following result can easily be obtained.

Proposition B.4 *Assume the hypotheses of Thm. B.3. Then*

$$f(a^w) - f(a)^w \in \Psi^w(M^p \lambda^{-1}, g),$$

where the function $\lambda(X)$ is defined in Subsect. B.1.

Note that the same result holds for the $(1, 0)$ quantization, thanks to (B.12).

Proof. one first proves the result for $f(\lambda) = (\lambda - z)^{-1}$, $z \in \mathbb{C} \setminus \mathbb{R}$, which amounts to construct a so-called parametrix for $a^w - z$. From symbolic calculus it follows that if $b_z(x, \xi) = (a(x, \xi) - z)^{-1}$, then $b_z^w(a^w - z) - \mathbb{1} \in \Psi^w(\lambda^{-1}, g)$. To extend the result to arbitrary functions one expresses $f(a^w)$ in terms of $(a^w - z)^{-1}$ using the well known functional calculus formula based on an almost analytic extension of f (see eg [DG1, Prop. C.2.2]). \square

B.4 Various estimates

The following lemma is proved in [A, Lemma 3.3].

Lemma B.5 *For $s \in [0, 1]$, and $v_i \in B(\mathcal{K}, \mathcal{K} \otimes \mathfrak{h})$, $i = 1, 2$ we have*

- 1) $\|(N + 1)^{-\frac{s}{2}} a(v_1) (H_0 + 1)^{-\frac{1-s}{2}}\| \leq \|\omega^{\frac{s-1}{2}} v_1\|_{B(\mathcal{K}, \mathcal{K} \otimes \mathfrak{h})}$,
- 2) $\|(H_0 + 1)^{-\frac{s}{2}} a^*(v_1) (N + 1)^{-\frac{1-s}{2}}\| \leq \|\omega^{-\frac{s}{2}} v_1\|_{B(\mathcal{K}, \mathcal{K} \otimes \mathfrak{h})}$,
- 3) $\|(N + 1)^{-s} a(v_1) a(v_2) (H_0 + 1)^{-1+s}\| \leq \|\omega^{-\frac{1-s}{2}} v_1\|_{B(\mathcal{K}, \mathcal{K} \otimes \mathfrak{h})} \|\omega^{-\frac{1-s}{2}} v_2\|_{B(\mathcal{K}, \mathcal{K} \otimes \mathfrak{h})}$,
- 4) $\|(H_0 + 1)^{-s} a^*(v_1) a^*(v_2) (N + 1)^{-1+s}\| \leq \|\omega^{-\frac{s}{2}} v_1\|_{B(\mathcal{K}, \mathcal{K} \otimes \mathfrak{h})} \|\omega^{-\frac{s}{2}} v_2\|_{B(\mathcal{K}, \mathcal{K} \otimes \mathfrak{h})}$.

The following theorem follows from the KLMN theorem and [RS1, Thm. VIII.25].

Theorem B.6 *Let H_0 be a positive selfadjoint operator on a Hilbert space \mathcal{H} . Let for $\kappa < \infty$, B_κ be quadratic forms on $D(H_0^{\frac{1}{2}})$ such that*

$$|B_\kappa(\psi, \psi)| \leq a \|H_0^{\frac{1}{2}} \psi\|^2 + b \|\psi\|^2,$$

where $a < 1$ uniformly in κ and $B_\kappa \rightarrow B_\infty$ on $D(H_0^{\frac{1}{2}})$.

Then

- (1) *there exists a selfadjoint operators H_κ with $D(H_\kappa) \subset D(H_0^{\frac{1}{2}})$ and*

$$(H_\kappa \psi, \psi) = B_\kappa(\psi, \psi) + (H_0^{\frac{1}{2}} \psi, H_0^{\frac{1}{2}} \psi), \quad \psi \in D(H_\kappa) \text{ for } \kappa \leq \infty.$$

- (2) *the resolvent $(z - H_\kappa)^{-1}$ converges in norm to $(z - H_\infty)^{-1}$.*

- (3) *e^{itH_κ} converges strongly to e^{itH_∞} when $\kappa \rightarrow +\infty$.*

References

- [A] Ammari, Z.: Asymptotic completeness for a renormalized nonrelativistic Hamiltonian in quantum field theory: the Nelson model, *Math. Phys. Anal. Geom.* **3** (2000), 217-285.
- [AHH] Arai, A., Hirokawa, M., Hiroshima, F.: On the absence of eigenvectors of Hamiltonians in a class of massless quantum field models without infrared cutoff, *J. Funct. Anal.*, **168** (1999), 470-497.
- [BFS] Bach, V., Fröhlich, J., Sigal, I. M.: Quantum electrodynamics of confined non-relativistic particles, *Adv. Math.*, **137** (1998), 299-395 .
- [BHLMS] Betz, V., Hiroshima, F., Lörinczi, J., Minlos, R. A., Spohn, H., Ground state properties of the Nelson Hamiltonian – a Gibbs measure-based approach, *Rev. Math. Phys.*, **14** (2002), 173-198 .

- [Bo] Bony, J. M.: Caractérisations des opérateurs pseudodifférentiels, Séminaire EDP, Centre de Mathématiques Laurent Schwartz, 1996-1997.
- [BD] Bruneau, L., Dereziński, J.: Pauli-Fierz Hamiltonians defined as quadratic forms, Rep. Math. Phys. **54** (2004), 169-199.
- [D] Davies, E. B.: Heat kernels and spectral theory, Cambridge Tracts in Mathematics. **92**. Cambridge university press (1989).
- [DG1] Dereziński, J., Gérard, C.: *Scattering Theory of Classical and Quantum N..Particle Systems*, Texts and Monographs in Physics, Springer-Verlag (1997).
- [DG2] Dereziński, J., Gérard, C.: Asymptotic completeness in quantum field theory. Massive Pauli-Fierz Hamiltonians, Rev. Math. Phys. **11** (1999), 383-450.
- [DG3] Dereziński, J., Gérard, C.: Scattering theory of infrared divergent Pauli-Fierz Hamiltonians, Annales Henri Poincaré, **5** (2004), 523-578.
- [GGM] Georgescu, V., Gérard, C., Moeller, J.: Spectral theory of massless Nelson models, Comm. Math. Phys. **249** (2004), 29-78.
- [G] Gérard, C.: On the existence of ground states for massless Pauli-Fierz Hamiltonians. Ann. Henri Poincaré **1** (2000), p 443-45.
- [GHPS1] Gérard, C., Hiroshima, F., Panati, A., Suzuki, A.: Infrared Divergence of a Scalar Quantum Field Model on a Pseudo Riemannian Manifold, *Interdisciplinary Information Sciences* **15**, (2009) 399-421.
- [GHPS2] Gérard, C., Hiroshima, F., Panati, A., Suzuki, A.: in preparation.
- [GP] Gérard, C., Panati, A.: Spectral and scattering theory for some abstract QFT Hamiltonians, Rev. Math. Phys. **21** (2009), 373-437.
- [Gr] Griesemer, M. Exponential decay and ionization thresholds in non-relativistic quantum electrodynamics. J. Funct. Anal. **210** (2004), 321-340.
- [GLL] Griesemer, M., Lieb, E., Loss, M.: Ground states in non-relativistic quantum electrodynamics. Invent. Math. **145** (2001), 557-595.
- [H] Hirokawa, M.: Infrared catastrophe for Nelson's model-non-existence of ground state and soft-boson divergence, Publ. RIMS, Kyoto Univ., **42** (2006), 897-922.
- [LMS] Lörinczi, J., Minlos, R. A., and Spohn, H.: The infrared behaviour in Nelson's model of a quantum particle coupled to a massless scalar field, Ann. Henri Poincaré, **3**(2002), 1-28.
- [MS] Milman, P. D. and Semenov, Y. A.: Global heat kernel bounds via desingularizing weights, J. Funct. Anal. **212** (2004) 373-398.

- [Ne] Nelson, E.: Interaction of nonrelativistic particles with a quantized scalar field, *J. Math. Phys.* **5** (1964), 1190-1997.
- [P] Panati, A.: Existence and nonexistence of a ground state for the massless Nelson model under binding condition. *Rep. Math. Phys.* **63** (2009), 305-330.
- [PE] Porper, F.O., Eidel'man, S.D.: Two sided estimates of fundamental solutions of second order parabolic equations and some applications, *Russian Math. Surveys*, **39** (1984), 119-178.
- [RS1] Reed, M., Simon, B.: *Methods of Modern Mathematical Physics, I: Functional Analysis*, Academic Press (1975).
- [RS2] Reed, M., Simon, B.: *Methods of Modern Mathematical Physics, II: Fourier Analysis, Self-adjointness*, Academic Press (1975).
- [Se] Semenov, Y.A.: Stability of L^p -spectrum of generalized Schrödinger operators and equivalence of Green's functions, *IMRN* **12** (1997), 573-593 .
- [Si] Simon, B.: *Functional Integration and Quantum Physics*, Academic Press (1979).
- [Sp] Spohn, H.: Ground state of a quantum particle coupled to a scalar boson field, *Lett. Math. Phys.*, **44** (1998), 9-16.
- [Zh] Zhang, Q.S.: Large time behavior of Schroedinger heat kernels and applications, *Comm. Math. Phys.* **210** (2000), 371-398.